

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
2 August 2001 (02.08.2001)

PCT

(10) International Publication Number
WO 01/55111 A1

(51) International Patent Classification⁷: **C07D 211/26**,
243/08, 333/16, 333/56, 307/80, 307/85, A61K 31/155,
31/381, 31/343, 31/404, 31/4164, 31/445, 31/4025,
31/506, 31/501

[GB/GB]; Granta Park, Abington, Cambridge CB1 6GB
(GB). **BOWER, Justin, Fairfield** [GB/GB]; Granta Park,
Abington, Cambridge CB1 6GB (GB).

(21) International Application Number: PCT/GB01/00362

(74) Agents: **HOWARD, Paul, Nicholas et al.**; Carpmaels &
Ransford, 43 Bloomsbury Square, London WC1A 2RA
(GB).

(22) International Filing Date: 27 January 2001 (27.01.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0001926.5 27 January 2000 (27.01.2000) GB
60/178,433 27 January 2000 (27.01.2000) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(71) Applicant (*for all designated States except US*): **RIBO-
TARGETS LIMITED** [GB/GB]; Granta Park, Abington,
Cambridge CB1 6GB (GB).

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **DRYSDALE, Mar-
tin, James** [GB/GB]; Granta Park, Abington, Cambridge
CB1 6GB (GB). **STARKEY, Ian, David** [GB/GB]; Granta
Park, Abington, Cambridge CB1 6GB (GB). **SWAR-
BRICK, Terry, Mark** [GB/GB]; Granta Park, Abington,
Cambridge CB1 6GB (GB). **POTTER, Andrew, John**

Published:

— with international search report

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*



WO 01/55111 A1

(54) Title: BIARYL COMPOUNDS, THEIR PREPARATION AND THEIR USE IN THERAPY

(57) Abstract: The invention relates to biaryl compounds, their preparation and their use in the treatment of bacterial and viral infection.

BIARYL COMPOUNDS, THEIR PREPARATION AND THEIR USE IN THERAPY


The present invention relates to a class of chemical compounds, their preparation and their use in therapy, particularly in the treatment of viral and bacterial infection.

5

Although many pharmaceutical compounds and compositions are available for the treatment of viral and bacterial infections, there remains a continuing need for improved treatments.

- 10 The present inventors have discovered a new class of chemical compounds which are particularly useful in the treatment of viral and bacterial infection.

According to the present invention there is provided a compound of the formula



- 15 wherein

Ar is an aryl group,

X¹ is selected from O, S, SO, SO₂ and NR,

X² is selected from O, S, SO, SO₂, NR and CR₂,

X³ is CR₂,


- 20 Y¹ and Y² are independently selected from C₁-₁₂ alkylene, C₄-₁₂ arylene, C₄-₁₆ aralkylene, CO(C₁-₁₂ alkylene), CO(C₄-₁₂ arylene) and CO(C₄-₁₆ aralkylene) groups.

A and B are independently selected from groups comprising a group selected from:

- amine (-NR₂), amide (-CONR₂), amidine (-C(=NR)NR₂), thioamide (-CSNR₂), oxime (=NOR), hydroxylamine (-NHOR), hydroxamic acid (-CONROR), hydrazine (-NRNR₂), hydrazone (=NNR₂), sulphonamide (-SO₂NR₂), sulphinamide (-SONR₂), sulfoximine (-SO(=NR)-), urea (-NRCONR₂), guanidine (-NRC(=NR)NR₂), and aromatic and non-aromatic nitrogen heterocyclic groups.
- 25

each R is independently selected from H, C₁₋₁₂ alkyl and C₃₋₁₂ aryl, or any two R groups may together comprise a C₁₋₆ alkylene chain, and pharmaceutically acceptable derivatives thereof.

- 5 In the compounds of the present invention Ar is an aryl group as herein defined. Preferably, the aryl group is a monocyclic or fused polycyclic (preferably bicyclic such as [6,5], [6,6] and [5,5] systems) aromatic or heteroaromatic group. Aromatic groups include phenyl and naphthyl. Heteroaromatic groups are generally preferred to the corresponding aromatic group. Heteroaromatic groups may comprise one or more heteroatoms.
- 10 Monocyclic heteroaromatic groups include pyridyl, pyrrolyl, furanyl, thienyl and thiazolyl. Heteroaromatic groups may be bonded to the rest of the molecule either via a ring carbon atom or via a ring heteroatom. Preferred fused bicyclic heteroaromatic groups include [6,5] (such as indolyl, indolinyl, benzofuranyl, benzothienyl), [6,6] (such as quinoliny, isoquinoliny, quinoxaliny) and [5,5] fused bicyclic heteroaromatic groups. [6,5] ring
- 15 systems, in which a heteroatom may be located at any ring position, are preferred. Particularly preferred fused bicyclic heteroaromatic groups comprise groups of the structure:



wherein X⁴ is NH, S or O. Indoles (i.e. where X⁴ is NH) are preferred.

20

Bicyclic heteroaromatic groups of this structure may be bonded to the rest of the molecule via any position, bonding via the 2, 3, 5 or 6 position being preferred.

The group Ar may be substituted as herein defined. Where substituted, there are preferably

25 one to three substituents, more preferably one substituent.

X¹ may be O, S, SO, SO₂ or NR. Preferably, X¹ is O.

X² may be O, S, SO, SO₂, NR or CR₂. Preferably, X² is NR, more preferably NH.

X^3 is CR_2 . Preferably, X^3 is CH_2 .

Y^1 and Y^2 are independently selected from C_{1-12} alkylene, C_{4-12} arylene, C_{4-16} aralkylene, $CO(C_{1-12}$ alkylene), $CO(C_{4-12}$ arylene) and $CO(C_{4-16}$ aralkylene) groups, as herein defined.

5

Preferably, Y^1 comprises a direct chain of 1 to 5 carbon atoms linking X^1 and A. For example, if Y^1 is an ethylene or *o*-phenylene group, the direct chain linking X^1 and A has two carbon atoms. Preferably, Y^1 comprises a C_{1-5} alkylene group.

- 10 Preferably, Y^2 comprises a direct chain of 1 to 5 carbon atoms linking X^2 and B. Preferably, Y^2 comprises a C_{1-5} alkylene group.

A and B are independently selected from groups comprising a group selected from amine ($-NR_2$), amide ($-CONR_2$), amidine ($-C(=NR)NR_2$), thioamide ($-CSNR_2$), oxime ($=NOR$),


- 15 hydroxylamine ($-NHOR$), hydroxamic acid ($-CONROR$), hydrazine ($-NRNR_2$), hydrazone ($=NNR_2$), sulphonamide ($-SO_2NR_2$), sulphinamide ($-SONR_2$), sulfoximine ($-SO(=NR)-$), urea ($-NRCONR_2$), guanidine ($-NRC(=NR)NR_2$), and aromatic and non-aromatic nitrogen heterocyclic groups.

- 20 Preferably, A and B are independently selected from groups comprising a group selected from amine, amidine, guanidine, and aromatic and non-aromatic nitrogen heterocyclic groups. Preferably, the amine, amidine and guanidine groups are unsubstituted (i.e. $R=H$).


The aromatic and non-aromatic nitrogen heterocyclic groups may be monocyclic
 25 (preferably 5 or 6 membered rings) or polycyclic (preferably fused bicyclic, more preferably [6,5], [6,6] and [5,5] systems) and may comprise one or more nitrogen atom. Examples of aromatic nitrogen heterocyclic groups include pyrrolyl, pyridinyl, 2-,3- and 4-pyrimidinyl, quinolinyl, isoquinolinyl, indolinyl, benzodiazolyl, benzotriazolyl, imidazolyl, triazolyl and thiazolyl groups. Examples of non-aromatic nitrogen heterocyclic
 30 groups include pyrrolidinyl, pyrrolidinone, piperidinyl, morpholinyl and piperazinyl groups. The aromatic and non-aromatic nitrogen heterocyclic groups may be substituted or unsubstituted. Preferred substituents include amino groups ($-NR_2$). The aromatic and non-

aromatic nitrogen heterocyclic group may be bonded to the rest of the molecule via a ring carbon atom or via a ring nitrogen atom or via a substituent.

- Included within the scope of the term aromatic and non-aromatic nitrogen heterocyclic groups are cyclic groups which mimic amidine or guanidine groups of the general formulae



- Specific examples include 2-aminopyridine, 2-aminopyrimidine and 2-pyrimidine groups:



- Each R is independently selected from H, C₁₋₁₂ alkyl and C₃₋₁₂ aryl, or any two R groups may together comprise a C₁₋₆ alkylene chain. For example, an R group in X² may be combined with an R group in B such that together with Y² a cyclic link is formed between X² and B.

- As used herein, the term "alkyl" means a branched or unbranched, cyclic or acyclic, saturated or unsaturated (e.g. alkenyl or alkynyl) hydrocarbyl radical. Where acyclic, the alkyl group is preferably a C₁₋₁₂, more preferably C₁₋₄ chain. Where cyclic, the alkyl group is preferably a C₃₋₁₂, more preferably C₅₋₁₀ and more preferably comprises a C₅, C₆ or C₇ ring. The alkyl chain or ring may include (i.e. be optionally interrupted with and/or terminate with) one or more heteroatoms, such as oxygen, sulphur or nitrogen.

- As used herein the term "alkylene" means a branched or unbranched, cyclic or acyclic, saturated or unsaturated divalent hydrocarbyl radical. Where acyclic the alkylene group is preferably a C₁₋₁₂, more preferably C₁₋₅ chain. Where cyclic, the alkylene group is

preferably a C₃₋₁₂, more preferably C₅₋₁₀, more preferably comprises a C₅, C₆ or C₇ ring. The alkylene chain or ring may include (i.e. be interrupted and/or terminate with) one or more heteroatoms such as oxygen, sulfur or nitrogen.

- 5 As used herein, the term "aryl" means a C₃₋₂₆, preferably C₃₋₁₂, aromatic group, such as phenyl or naphthyl, or a heteroaromatic group containing one or more, preferably one, heteroatom, such as pyridyl, pyrrolyl, furanyl, thienyl, thiazolyl, indolyl, indolinyl, benzofuranyl, benzothienyl, quinolinyl, isoquinolinyl, quinoxaliny, 2-, 3- or 4-pyrimidinyl, benzodiazolyl, benzotriazolyl, imidazolyl, triazolyl and thiazolyl groups.

10

As used herein the term "arylene" means a divalent hydrocarbyl radical comprising a C₃₋₁₂ aromatic group (such as *o*-, *m*- or *p*-phenylene) or heteroaromatic group containing one or more, preferably one, heteroatom (such as a pyridine-2,3-diyl group).

- 15 As used herein the term "aralkylene" means a divalent hydrocarbyl radical comprising both alkylene and arylene groups (such as -CH₂-(*o*-phenylene)-CH₂-).

- The alkyl, aryl, alkylene, arylene and aralkylene groups Ar, Y¹, Y² and R, and the groups A and B, may be further substituted or unsubstituted. For example, a C₁ (methyl) group may be
- 20 further substituted with a phenyl group to give a benzyl group. Substituents may include carbon containing groups such as alkyl, aryl, aralkyl (e.g. substituted and unsubstituted phenyl, substituted and unsubstituted benzyl); halogen atoms (e.g. F, Cl, Br and I) and halogen containing groups such as haloalkyl (e.g. trifluoromethyl); oxygen containing groups such as alcohols (e.g. hydroxy, hydroxyalkyl, aryl(hydroxy)alkyl), ethers (e.g.
- 25 alkoxy, alkoxyalkyl, aryloxyalkyl), aldehydes (e.g. carboxaldehyde), ketones (e.g. alkylcarbonyl, alkylcarbonylalkyl, arylcarbonyl, arylalkylcarbonyl, arylcarbonylalkyl), acids (e.g. carboxy, carboxyalkyl), acid derivatives such as esters (e.g. alkoxycarbonyl, alkoxycarbonylalkyl, alkylcarbonyloxy, alkylcarbonyloxyalkyl) and amides (e.g. aminocarbonyl, mono- or dialkylaminocarbonyl, aminocarbonylalkyl, mono- or
- 30 dialkylaminocarbonylalkyl, arylaminocarbonyl); and carbamates (eg. alkoxycarbonylamino, aryloxy carbonylamino, aminocarbonyloxy, mono- or dialkylaminocarbonyloxy, arylaminocarbonyloxy), and ureas (eg. mono- or dialkylaminocarbonylamino or arylaminocarbonylamino); nitrogen containing groups such

as amines (e.g. amino, mono- or dialkylamino, aminoalkyl, mono- or dialkylaminoalkyl), azides, nitriles (e.g. cyano, cyanoalkyl), nitro; sulfur containing groups such as thiols, thioethers, sulfoxides, and sulfones (e.g. alkylthio, alkylsulfinyl, alkylsulfonyl, alkylthioalkyl, alkylsulfinylalkyl, alkylsulfonylalkyl, arylthio, arylsulfinyl, arylsulfonyl, 5 arylthioalkyl, arylsulfinylalkyl, arylsulfonylalkyl); and heterocyclic groups containing one or more, preferably one, heteroatom (e.g. thienyl, furanyl, pyrrolyl, imidazolyl, pyrazolyl, thiazolyl, isothiazolyl, oxazolyl, oxadiazolyl, thiadiazolyl, pyrrolidinyl, pyrrolinyl, imidazolidinyl, imidazoliny, pyrazolidinyl, tetrahydrofuranyl, pyranal, pyronyl, pyridyl, pyrazinyl, pyridazinyl, piperidyl, piperazinyl, morpholinyl, thianaphthyl, benzofuranyl, 10 isobenzofuranyl, indolyl, oxyindolyl, isoindolyl, indazolyl, indoliny, 7-azaindolyl, benzopyranal, coumarinyl, isocoumarinyl, quinoliny, isoquinoliny, naphthridiny, cinnoliny, quinazoliny, pyridopyridyl, benzoxazinyl, quinoxaliny, chromenyl, chromanyl, isochromanyl, phthalazinyl and carboliny).

15 As used herein, the term "alkoxy" means alkyl-O- and "alkanoyl" means alkyl-CO. Alkyl substituent groups or alkyl-containing substituent groups may comprise one or more further substituents. As used herein, the term "aryloxy" means aryl -O- and "aryloyl" means aryl -CO. Aryl substituent groups or aryl-containing substituent groups may comprise one or more further substituents.

20

As used herein, the term "halogen" means a fluorine, chlorine, bromine or iodine radical, preferably a fluorine or chlorine radical.

By "a pharmaceutically acceptable derivative" is meant any pharmaceutically acceptable 25 salt, addition compound, or any other compound which upon administration to a recipient is capable of providing (directly or indirectly) a compound of the present invention or a pharmaceutically acceptable metabolite. By "pharmaceutically acceptable metabolite" is meant a metabolite or residue of a compound of the present invention which gives rise to a biological activity exhibited by the present compounds.

30

As used herein, a "patient" is a mammal (e.g., such as a human being or other non-human mammal) to whom a compound according to the invention is administered. The term

"patient" does not imply that the individual has ever been hospitalized for medical treatment.

As used herein, a "microorganism" refers to a bacterial, viral, prokaryotic or eukaryotic organism which can be viewed microscopically. The term "microorganism" as used herein
5 encompasses both bacteria and viruses.

As used herein, "anti-microbial properties" or "anti-bacterial properties" or "anti-viral properties" refer to the ability of the compounds according to the invention to inhibit microbial, bacterial, viral growth. As defined herein, "inhibiting growth" refers to an inhibition in the translation of microbial proteins, which in turn results in an inhibition in
10 microbial replication (and therefore transcription of microbial mRNAs) which in turn results in an inhibition of infection. Any one of these processes (e.g., translation, replication, transcription, infection) may be assayed to determine the effectiveness of the compounds according to the invention (e.g., defined as the ability of the compound to inhibit growth). As defined herein, "inhibition of microbial growth" refers to an at least
15 two-fold decrease in any of the parameters discussed above (e.g., translation of microbial proteins, replication of microorganisms, transcription of microbial mRNAs, and/or infection by microorganisms). Inhibition can also refer to an at least two fold decrease in an immune response associated with a microbial infection (e.g., such as the accumulation of anti-microbial antibodies or cytokines and/or pyrogens associated with microbial
20 infection). In one embodiment, inhibition is at least 2-fold, at least 10-fold, at least 20-fold, at least 30-fold, at least 40-fold, at least 50-fold, or at least 100-fold.

In a preferred embodiment of the invention, the compound inhibits translation of a bacterial and/or viral transcript. In still a more preferred embodiment, the compound inhibits translation of a bacterial and/or viral transcript while not inhibiting translation of a
25 mammalian transcript. In one embodiment of the invention, translation is inhibited at least 2-fold, at least 5-fold, at least 10-fold, at least 20-fold, at least 30-fold, at least 40-fold, at least 50-fold, or at least 100-fold compared to translation of bacterial and/or viral transcripts in a mammalian organism which has not been treated with the compounds according to the invention. In a further embodiment, the compound inhibits bacterial
30 and/or viral replication.

According to a further aspect of the present invention there is provided a compound according to the present invention for use in a method of treatment, preferably in the prophylaxis or treatment of viral infection or bacterial infection.

- 5 According to a further aspect of the present invention there is provided use of a compound according to the present invention in the manufacture of a medicament for the prophylaxis or treatment of viral infection or bacterial infection.

10 According to a further aspect of the present invention there is provided a method of prophylaxis or treatment of viral infection or bacterial infection comprising administration to a patient in need of such treatment an effective dose of a compound according to the present invention.

In one embodiment, the effective dose of the compound according to the invention is a
15 dose effective to decrease the titer of infectious microorganisms in a patient's body. In one embodiment, the titer of infectious microorganisms is measured by culturing a bodily sample and counting the number of microorganisms in the sample. In another embodiment, the titer of infectious microorganisms is determined by measuring the expression of the bacterial or viral nucleic acids and/or proteins. In a further embodiment,
20 the effective dose of the compound is a dose effective to restore the immune response of a host (e.g., a patient) to a microorganism to normal (e.g., to resemble an immune response of an uninfected host). For example, in one embodiment, a bodily fluid from a patient is assayed to detect the presence and/or amounts of anti-bacterial or antiviral antibodies.

In one embodiment, a compound according to the invention is administered to a patient
25 who has both a bacterial and a viral infection. In one embodiment, the patient treated has AIDS. In another embodiment, the person has AIDS and at least one opportunistic infection.

In another embodiment, a compound according to the invention is used prophylactically. In one embodiment, the compound is contacted with a cell or surface thereby to prevent the
30 growth of microorganisms in proximity to the cell or surface. In one embodiment, the compound is administered to a patient to prevent infection by a microorganism or to reduce

the severity of infection (e.g., as measured by determining the titer of the microorganism in a treated vs. an untreated individual).

Viral infections include, but are not limited to:

5			
	<i>Family</i>	<i>Virus</i>	<i>Disease</i>
	<u>Adenoviruses</u>	Adenovirus	acute respiratory disease
	<u>Arenaviruses</u>	Lassa Virus	Lassa Fever
	<u>Astroviridae</u>	Astrovirus	Enteritis
10	<u>Bunyaviridae</u>	Hantavirus Phlebovirus	Hantavirus Pulmonary Syndrome Rift Valley Fever
	<u>Calciviridae</u>	Hepatitis E	
	<u>Filoviridae</u>	Ebola Virus Marburg Virus	
15	<u>Flaviviridae</u>	Japanese Encephalitis Virus Hepatitis C Virus Dengue Virus Yellow Fever Virus Hepatitis G Virus	Dengue Haemorrhagic Fever
20	<u>Hepadnaviridae</u>	Hepatitis B Virus Hepatitis D (delta) Virus	
	<u>Herpesviridae</u>	Herpes Simplex Virus 1 Herpes Simplex Virus 2 Cytomegalovirus (CMV) Epstein Barr Virus (EBV) Varicello Zoster Virus (VZV) HHV-6 HHV-7 KSHV/HHV8	Mononucleosis Chicken Pox/Shingles Kaposi Sarcoma
25			
30	<u>Orthomyxoviruses</u>	Influenza Virus	
	<u>Paramyxoviridae</u>	Paramyxoviruses Rubulaviruses Morbilliviruses Respiratory Syncytial Virus	Para-Influenza Mumps Measles
35	<u>Papovaviridae</u>	Papillomaviruses Polyomaviruses BK and JC Virus	Warts/Cervical Cancer
	<u>Parvoviridae</u>	Parvoviruses	Erythema Infectiosum
40	<u>Picomaviridae</u>	Coxsackie Viruses (A and B)	Viral Myocarditis & Meningitis & Enteritis

	Hepatitis A Virus	Hepatitis
	Polioviruses	Poliomyelitis
	Rhinoviruses	Cold
5	<u>Reoviridae</u>	Astroviruses
		Caliciviruses
		Reoviruses
	<u>Rhabdoviridae</u>	Lyssavirus
	<u>Retroviridae</u>	HIV-1 and HIV-2
10		HTLV-1 and HTLV-2
		Diarrhoea
		Diarrhoea
		Diarrhoea
		Rabies
		AIDS
		Leukaemia

Preferably, the viral infection comprises HIV or HCV infection, more preferably HIV-I or HIV-II.

15

Bacterial infections include, but are not limited to, infections by *Gram Positive Bacteria* including *Bacillus cereus*, *Bacillus anthracis*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium tetani*, *Clostridial perfringens*, *Corynebacteria diphtheriae*, *Enterococcus* (*Streptococcus D*), *Listeria Monocytogenes*, *Pneumococcal Infections* (*Streptococcus pneumoniae*), *Staphylococcal Infections* and *Streptococcal Infections*; *Gram Negative Bacteria* including *Bactroides*, *Bordetella pertussis*, *Brucella*, *Campylobacter Infections*, *Enterohemorrhagic Escherichia coli* (*EHEC/E.coli O157:H7*), *Enteroinvasive Escherichia coli* (*EIEC*), *Enterotoxigenic Escherichia coli* (*ETEC*), *Haemophilus influenzae*, *Helicobacter pylori*, *Klebsiella pneumonia*, *Legionella spp.*, *Moraxella catarrhalis*, *Neisseria gonorrhoeae*, *Neisseria meningitidis*, *Proteus spp.*, *Pseudomonas aeruginosa*, *Salmonella spp.*, *Shigella spp.*, *Vibrio cholera* and *Yersinia*; *Acid Fast Bacteria* including *Mycobacterium tuberculosis*, *Mycobacterium avium-intracellulare*, *Mycobacterium leprae*, *Atypical Bacteria*, *Chlamydia*, *Mycoplasma*, *Rickettsia*, *Spirochetes*, *Treponema pallidum*, *Borrelia recurrentis*, *Borrelia burgdorferi* and *Leptospira* *icterohemorrhagiae*; and other miscellaneous bacteria including *Actinomyces* and *Nocardia*.

It is a feature of the compounds of the present invention that they inhibit the binding of the HIV protein Tat to the HIV RNA Tar binding site. Accordingly, the present invention further provides use of a compound of the present invention to inhibit the binding of Tat to Tar.

35


It is also feature of the compounds of the present invention that they inhibit translation of bacterial proteins. Accordingly, the present invention further provides use of a compound of the present invention to inhibit the translation of bacterial proteins.

- 5 According to a further aspect of the present invention there is provided a pharmaceutical composition comprising a compound of the present invention in combination with a pharmaceutically acceptable excipient.

- 10 According to a further aspect of the present invention there is provided a method of preparing a pharmaceutical composition comprising the step of combining a compound of the present invention with a pharmaceutically acceptable excipient.

- According to a further aspect of the present invention there is provided a process for the preparation of the compounds of the present invention. The compounds of the present
15 invention may be prepared according to the following general reaction scheme.

General Reaction Scheme



Reagents (i) R^1 halide, Cs_2CO_3 ; (ii) $ArB(OH)_2$, Pd catalyst; (iii) R^2R^3NH , DCE or EtOH, reducing agent; (iv) $R^2CH_2Br^- Ph_3P^+$, base, toluene; (v) H_2 , Pd/C, solvent; (vi) (a) $NaBH_4$, solvent, (b) Ph_3P , CBr_4 ; (vii) HZ^2R^2 , base, solvent.

- 5 The biaryl compounds according to the invention have anti-microbial (e.g., anti-bacterial and/or anti-viral properties). In one embodiment, the compounds inhibit microbial growth. Inhibition of microbial growth can be assayed in a number of different ways. In one embodiment, microbial growth is measured by assaying the translation of microbial proteins, levels of microbial replication, transcription of microbial mRNAs, and infectivity
- 10 (e.g., viral titer in cells exposed to a virus). Assays for measuring such parameters are well known in the art and include, but are not limited to, immunossays to detect translation products or assays which measure binding of translational regulators to mRNA transcripts (e.g., to measure translation), RT-PCT, or hybridization assays (e.g., to measure transcription), incorporation of labeled nucleotides or hybridization assays to measure the
- 15 presence/amount of microbial genomic DNA (e.g., to measure replication), plate counting assays (e.g., to measure microbial titers), and the like.

- In one embodiment, compounds are synthesized according to the methods described above and the ability of the compounds to inhibition of microbial growth is assayed to identify compounds which produce an at least two-fold decrease in any of the parameters discussed
- 20 above (e.g., translation of microbial proteins, replication of microorganisms, transcription of microbial mRNAs, and/or infection by microorganisms). In one embodiment, inhibition is at least 2-fold, at least 10-fold, at least 20-fold, at least 30-fold, at least 40-fold, at least 50-fold, or at least 100-fold. In one embodiment, activity is measured *in vitro*, e.g., by measuring the effects of the compounds on bacterial cultures or on cells infected or to be
- 25 infected with a virus. In another embodiment, compounds are selected which inhibit the growth of both bacterial and viral microorganisms. In one embodiment, compounds are selected which inhibit the growth of HIV in cells infected or to be infected with the virus. In still another embodiment, compounds are selected which inhibit the growth of HIV and any of the characteristic microorganisms found in opportunistically infected AIDS patients.
- 30 In another embodiment, compounds according to the invention are tested in animal models to determine the effects of the compounds on microbial growth as described above. In one

embodiment, the compounds are tested for their affect on the immune response of an animal to a microbial infection to select compounds which return the immune response to normal (e.g., provide a response similar to that observed in an animal which has not been infected. For example, in one embodiment, a bodily fluid (e.g., blood) is obtained from an
5 infected animal at various time points after administering a compound according to the invention to determine the presence or absence of antibodies specific for microbial antigens and/or the presence or absence of cytokines characteristic of microbial infection. Additionally, or alternatively, the animal may be tested by evaluating any of the parameters discussed above (e.g., translation of microbial proteins, replication of
10 microorganisms, transcription of microbial mRNAs, and/or infection by microorganisms).

In a preferred embodiment, the compounds according to the invention inhibit translation of a bacterial and/or viral transcript. In still a more preferred embodiment, the compound inhibits translation of a bacterial and/or viral transcript while not inhibiting translation of a mammalian transcript. In one embodiment of the invention, translation is inhibited at least
15 2-fold, at least 5-fold, at least 10-fold, at least 20-fold, at least 30-fold, at least 40-fold, at least 50-fold, or at least 100-fold compared to translation of bacterial and/or viral transcripts in a mammalian organism which has not been treated with the compounds according to the invention. In one embodiment, a reporter gene is cloned downstream an in frame with a bacterial or viral translation initiation sequence. and the activity of the
20 compounds synthesised is assayed by monitoring the presence and/or amount of the protein encoded by the reporter gene.

In a preferred embodiment, a compound according to the invention is provided which inhibits the binding of the HIV protein Tat to the HIV RNA Tar binding site. Accordingly, the present invention further provides use of a compound of the present invention to inhibit
25 the binding of Tat to Tar. In one embodiment, inhibition is measured directly by measuring binding of Tat to Tar. In another embodiment, inhibition is measured by measuring the production of Tat protein.

In another embodiment, the compounds according to the invention are tested for their ability to prevent microbial infection. For example, in one embodiment, the compounds
30 are contacted to a cell and the ability of a microorganism to grow in proximity to said cell

is evaluated. In one embodiment, the cell is a cell which is to be infected with a virus. and the cell is contacted with the compound prior to contacting the cell with the virus. The ability of the compounds to be used prophylactically is then evaluated as described above (e.g., by assaying one or more of translation of microbial proteins, replication of
5 microorganisms. transcription of microbial mRNAs, and/or infection by microorganisms). In a further embodiment, the compounds according to the invention are contacted with a surface and assayed for their ability to prevent microbial growth on the surface.

The medicament employed in the present invention can be administered by oral or parenteral routes, including intravenous, intramuscular, intraperitoneal, subcutaneous.
10 transdermal, airway (aerosol), rectal, vaginal and topical (including buccal and sublingual) administration.

For oral administration, the compounds of the invention will generally be provided in the form of tablets or capsules, as a powder or granules, or as an aqueous solution or
15 suspension.

Tablets for oral use may include the active ingredients mixed with pharmaceutically acceptable excipients such as inert diluents, disintegrating agents, binding agents, lubricating agents, sweetening agents, flavouring agents, colouring agents and
20 preservatives. Suitable inert diluents include sodium and calcium carbonate, sodium and calcium phosphate, and lactose, while corn starch and alginic acid are suitable disintegrating agents. Binding agents may include starch and gelatin, while the lubricating agent, if present, will generally be magnesium stearate, stearic acid or talc. If desired, the tablets may be coated with a material such as glyceryl monostearate or glyceryl distearate,
25 to delay absorption in the gastrointestinal tract.

Capsules for oral use include hard gelatin capsules in which the active ingredient is mixed with a solid diluent, and soft gelatin capsules wherein the active ingredients is mixed with water or an oil such as peanut oil, liquid paraffin or olive oil.

30

Formulations for rectal administration may be presented as a suppository with a suitable base comprising for example cocoa butter or a salicylate.

Formulations suitable for vaginal administration may be presented as pessaries, tampons, creams, gels, pastes, foams or spray formulations containing in addition to the active ingredient such carriers as are known in the art to be appropriate.

5

For intramuscular, intraperitoneal, subcutaneous and intravenous use, the compounds of the invention will generally be provided in sterile aqueous solutions or suspensions, buffered to an appropriate pH and isotonicity. Suitable aqueous vehicles include Ringer's solution and isotonic sodium chloride. Aqueous suspensions according to the invention
10 may include suspending agents such as cellulose derivatives, sodium alginate, polyvinylpyrrolidone and gum tragacanth, and a wetting agent such as lecithin. Suitable preservatives for aqueous suspensions include ethyl and n-propyl p-hydroxybenzoate.

The compounds of the invention may also be presented as liposome formulations.

15

In general a suitable dose will be in the range of 0.01 to 100 mg per kilogram body weight of the recipient per day, preferably in the range of 0.2 to 10 mg per kilogram body weight per day. The desired dose is preferably presented once daily, but may be dosed as two, three, four, five or six or more sub-doses administered at appropriate intervals throughout
20 the day. These sub-doses may be administered in unit dosage forms, for example, containing 10 to 1500 mg, preferably 20 to 1000 mg, and most preferably 50 to 700 mg of active ingredient per unit dosage form.

In a preferred method according to the present invention, the compounds are used to
25 prevent or delay the onset of HIV-infection in individuals who are susceptible or at risk of HIV-infection (e.g., intravenous drug users, patients who have had, or are about to receive, a blood transfusion, immunodeficient or immunocompromised patients, gay men, and the like). The method comprises administering to such a patient a prophylactically effective amount (which generally is the same as a therapeutically effective amount) of one or more
30 of the compositions according to the present invention to the patient to delay or prevent an HIV-infection. In another embodiment of the invention, the compounds are used to treat an already-infected patient (e.g., an HIV-positive patient) to prevent re-infection and to inhibit viral replication and/or further infection by opportunistic microorganisms. The

compounds may be used by themselves or in conjunction with other drugs (e.g., protease inhibitors, antibiotics) or other therapies.

The invention will now be described with reference to the following Examples. It will be appreciated that what follows is by way of example only and that modifications to detail may be made whilst still falling within the scope of the invention.

EXPERIMENTAL

10 Chemical Synthesis

The compounds of the present invention were synthesized according to the following protocols and characterized by standard spectroscopic techniques including LCMS under the following conditions.

15 HPLC: HP1100

Column: ABZ+, 3.3cm*4.6mmD

Temperature: 20°C

Solvents: A - Water + 0.1% formic acid + 10mmol ammonium acetate

B - 95% Acetonitrile/water + 0.05% formic acid

20 Flow rate: 1mL/min

Gradient: Total time 8 minutes

- 100% A for 0.7 minutes

- ramp up to 100% B over 3.5 minutes

- 100% B for 3.5 minutes


25

- ramp down to 0% B over 0.3 minutes

Detection: UV detection at 230nm, 254nm and 270nm

Mass spec: HP1100 MSD

Method: Electrospray, +ve ion




Reagents: (i) $\text{BrCH}_2\text{CH}_2\text{NHBoc}'$, Cs_2CO_3 , DMF; (ii) ArB(OH)_2 , $\text{PdCl}_2(\text{PPh}_3)_2$, DME, Na_2CO_3 ; (iii) $\text{BocNH(CH}_2)_4\text{NH}_2$, DCE, sodium triacetoxyborohydride; (iv) TFA/DCM, 1/1; (v) N,N' -bis-*t*-butoxycarbonylpyrazolecarboxamidine, N,N -diisopropylethylamine, CH_3CN .

Scheme 1


The following examples were synthesized using the procedures outlined in Scheme 1.

Example 1




- 15 To 5-bromosalicylaldehyde (1eq.) and cesium carbonate (2 eq.) in DMF at RT was added 1-bromo-2-*N*-*t*-butoxycarbonylethane (1.2 eq.), and the mixture stirred overnight at RT. The DMF was evaporated *in vacuo* and the residue partitioned between EtOAc and water. The organic layer was washed with brine and dried over MgSO_4 . Concentration gave 1 as a solid, which was crystallized from hexane.
- 20 LC retention time 4.56 minutes. $[\text{M}+\text{H}]^+$ 345.

Example 2



5

a)




To a mixture of the aldehyde of example 1 (1 eq.), dichlorobis(triphenylphosphine)-
 10 palladium(II) (10mol%) and 2N Na₂CO₃ (eq.) was added phenylboronic acid (1.5eq.) in
 dry, degassed DME and the mixture heated at 80°C for 16h. After cooling to RT the
 solvent was removed *in vacuo* and the residue partitioned between ethyl acetate and water.
 The organic layer was washed with brine and dried over MgSO₄. Concentration gave a
 solid, which was purified by chromatography on silica gel eluting with mixtures of ethyl
 15 acetate and hexane.

LC retention time 4.75 minutes, [M+Na]⁺ 364.

b)


20



The aldehyde (1 eq.) and mono-*N*-*t*-butoxycarbonyl-1,4-diaminobutane (2 eq.) were stirred
 at RT for 15min in 1,2-dichloroethane, and then sodium triacetoxyborohydride (1.5 eq.)
 was added. After stirring for 16h at RT the mixture was concentrated *in vacuo* and the
 25 residue partitioned between dichloromethane and water. The organic layer was washed
 with brine and dried over MgSO₄. Concentration gave a solid, which was purified by
 chromatography on silica gel eluting with mixtures of dichloromethane and methanol.

LC retention time 3.79 minutes, $[M+H]^+$ 514.

c)



The bis-Boc compound was treated with 1mL of 1/1 DCM/TFA and stirred at RT for 1h. The volatiles were removed *in vacuo* and the TFA salt used without further purification. The TFA salt was stirred in acetonitrile, treated with excess *N,N*-diisopropylethylamine and *N,N'*-bis-*t*-butoxycarbonylpyrazole carboxamidine added. The mixture was stirred overnight at RT then concentrated *in vacuo*. The residue was partitioned between dichloromethane and water, the organic layer washed with brine and dried over $MgSO_4$. Concentration gave an oil, which was purified by chromatography on silica gel eluting with mixtures of dichloromethane and methanol.


10

15 LC retention time 4.44 minutes, $[M+H]^+$ 799.

d) The fully protected bis-guanidine was treated with 1mL of 1/1 DCM/TFA and stirred at RT for 1h. The volatiles were removed *in vacuo* to give the desired bis-guanidine as the tris-trifluoroacetate.

20 LC retention time 0.53 and 2.64 minutes, $[M+H]^+$ 398.


Example 3



As example 2 using *p*-tolylboronic acid.


LC retention time 2.78 minutes, $[M+H]^+$ 412.

Example 4




- 5 As example 2 using 4-methoxyphenylboronic acid.
LC retention time 2.71 minutes, $[M+H]^+$ 428.

Example 5




- As example 2 using 4-fluorophenylboronic acid.
LC retention time 2.71 minutes, $[M+H]^+$ 416.

15 Example 6



- As example 2 using 2-thienylboronic acid.
- 20 LC retention time 0.50 and 2.62 minutes, $[M+H]^+$ 404.

Example 7




As example 2 using 3-thienylboronic acid.

LC retention time 0.51 and 2.58 minutes, $[M+H]^+$ 404.

5


Example 8



10 As example 2 using benzo[b]thiophene-2-boronic acid.

LC retention time 2.92 minutes, $[M+H]^+$ 454.


Example 9



15

As example 2 using benzo[b]furan-2-boronic acid.

LC retention time 2.85 minutes, $[M+H]^+$ 438.




Reagents: (i) Br(CH₂)₃NHBoc^t, Cs₂CO₃, DMF; (ii) ArB(OH)₂, PdCl₂(PPh₃)₂, DME, Na₂CO₃; (iii) (*N,N'*-bis-*t*-butoxycarbonylcarboxamidine)NH(CH₂)₄NH₂, DCE, sodium triacetoxyborohydride; (iv) TFA/DCM, 1/1.

5

Scheme 2

The following examples were synthesized using the procedures outlined in Scheme 2.

10 Example 10




To 5-bromosalicylaldehyde (1 eq) and cesium carbonate (2 eq) in DMF at RT was added 1-bromo-3-*N*-*t*-butoxycarbonylpropane (1.2) and the mixture stirred overnight at RT. The DMF was evaporated *in vacuo* and the residue partitioned between EtOAc and water. The organic layer was washed with brine and dried over MgSO₄. Concentration gave the desired aldehyde as a solid, which was crystallized from hexane.


LC retention time 4.68 minutes. [M+H]⁺ 380.

20

Example 11




5 a)



To a mixture of the aldehyde (1 eq) of example 10, dichlorobis(triphenylphosphine)-
 10 palladium(II) (10 mol%) and 2N Na₂CO₃ was added *p*-tolylboronic acid (2 eq) in dry, degassed DME and the mixture heated at 80°C for 16h. After cooling to RT the solvent was removed *in vacuo* and the residue partitioned between ethyl acetate and water. The organic layer was washed with brine and dried over MgSO₄. Concentration gave a solid,
 15 and hexane.

LC retention time 4.79 minutes, [M+H-Boc]⁺ 286.

b)



20

The aldehyde (1 eq) and 1-amino-4-*N,N'*-bis-*t*-butoxycarbonylguanidinobutane (1.8 eq) were stirred at RT for 15min in 1,2-dichloroethane, then sodium triacetoxyborohydride (1.5 eq) was added in one portion. After stirring for 16h at RT the mixture was
 25 concentrated *in vacuo* and the residue partitioned between dichloromethane and water. The organic layer was washed with brine and dried over MgSO₄. Concentration gave a solid, which was purified by chromatography on silica gel eluting with mixtures of dichloromethane and methanol.


LC retention time 4.18 minutes, $[M+H]^+$ 700.

- c) The tris-Boc compound was treated with 1mL of 1/1 DCM/TFA and stirred at RT for 1h. The volatiles were removed *in vacuo* to give the desired mono-guanidine as the tris-trifluoroacetate.

LC retention time 2.69 minutes, $[M+H]^+$ 400.

10

Example 12




15

As example 11 using 4-fluorophenylboronic acid.

LC retention time 2.68 minutes, $[M+H]^+$ 388.

Example 13

20




As example 11 using 3-thienylboronic acid.

LC retention time 0.49 and 2.56 minutes, $[M+H]^+$ 376.

25


Example 14



As example 11 using benzo[b]thiophene-2-boronic acid.

LC retention time 2.91 minutes, $[M+H]^+$ 426.


5 Example 15



As example 11 using benzo[b]furan-2-boronic acid.

10 LC retention time 2.84 minutes, $[M+H]^+$ 410.

Example 16




15

As example 11 using 4-trifluoromethylbenzeneboronic acid.

LC retention time 2.90 minutes, $[M+H]^+$ 438.


20 Example 17



As example 11 using naphthyl-1-boronic acid.

25 LC retention time 2.87 minutes, $[M+H]^+$ 420.


Example 18



5 As example 11 using naphthyl-2-boronic acid.


LC retention time 2.87 minutes, $[M+H]^+$ 420.

Example 19



As example 11 using 1-*N*-Boc-indole-2-boronic acid.

LC retention time 2.76 minutes, $[M+H]^+$ 410.




Reagents: (i) $\text{Br}(\text{CH}_2)_3\text{NHBoc}$, Cs_2CO_3 , DMF; (ii) $\text{ArB}(\text{OH})_2$, $\text{PdCl}_2(\text{PPh}_3)_2$, DME, Na_2CO_3 ; (iii) $\text{R}^1\text{R}^2\text{NH}_2$, DCE, sodium triacetoxyborohydride; (iv) TFA/DCM, 1/1.

Scheme 3

The following examples were synthesized using the procedures outlined in Scheme 3.

Example 20




- 5 To a mixture of the aldehyde (1 eq) of example 10, dichlorobis(triphenylphosphine)-palladium(II) (10 mol%) and 2N Na₂CO₃ was added benzo[*b*]furan-2-boronic acid acid (2 eq) in dry, degassed DME and the mixture heated at 80°C for 16h. After cooling to RT the solvent was removed *in vacuo* and the residue partitioned between ethyl acetate and water. The organic layer was washed with brine and dried over MgSO₄. Concentration gave a
- 10 solid, which was purified by chromatography on silica gel eluting with mixtures of ethyl acetate and hexane.

LC retention time 5.17 minutes, [M+Na]⁺ 418.


Example 21

15



a)

20



- The aldehyde (1 eq.) from example 20 and mono-*N,N'*-bis-*t*-butoxycarbonylguanidino-*m*-xylenediamine (2 eq.) were stirred at RT for 15min in 1,2-dichloroethane, then sodium triacetoxyborohydride (1.5 eq.) was added in one portion. After stirring for 16h at RT the
- 25 mixture was concentrated *in vacuo* and the residue partitioned between dichloromethane and water. The organic layer was washed with brine and dried over MgSO₄.


Concentration gave a solid, which was purified by chromatography on silica gel eluting with mixtures of dichloromethane and methanol.

LC retention time 4.38 minutes, $[M+H]^+$ 758.

- 5 b) The tris-Boc compound was treated with 1mL of 1/1 DCM/TFA and stirred at RT for 1h. The volatiles were removed *in vacuo* to give the desired mono-guanidine as the tris-trifluoroacetate.

LC retention time 2.90 minutes, $[M+H]^+$ 458.


10 Example 22



As example 21 using 4-*N,N'*-bis-*t*-butoxycarbonylguanidinoaminomethylaniline.

- 15 LC retention time 2.82 minutes, $[M+H]^+$ 444.


Example 23.



As example 21 using 1-amino-4-[(*N,N'*-bis-*t*-butoxycarbonyl)-*N*-methyl] guanidinobutane.

- 20 LC retention time 2.78 minutes, $[M+H]^+$ 424.

Example 24




25

As example 21 using mono-*N*-*t*-butoxycarbonyldiaminobutane.

LC retention time 2.74 minutes. $[M+H]^+$ 368.

Example 25




- 5 To a mixture of the aldehyde of example 10 (1 eq.), dichlorobis(triphenylphosphine)-palladium(II) (10 mol%) and 2N Na₂CO₃ was added benzo[*b*]thiophene-2-boronic acid (2 eq) in dry, degassed DME and the mixture heated at 80°C for 16h. After cooling to RT the solvent was removed *in vacuo* and the residue partitioned between ethyl acetate and water. The organic layer was washed with brine and dried over MgSO₄. Concentration gave a
- 10 solid, which was purified by chromatography on silica gel eluting with mixtures of ethyl acetate and hexane.


LC retention time 5.30 minutes, [M+H-Boc]⁺ 312.

Example 26

15



a)



20

- The aldehyde (1 eq.) from Example 25 and mono-*N,N'*-bis-*t*-butoxycarbonylguanidino-*m*-xylenediamine (2 eq.) were stirred at RT for 15min in 1,2-dichloroethane, then sodium triacetoxyborohydride (1.5 eq.) was added in one portion. After stirring for 16h at RT the
- 25 mixture was concentrated *in vacuo* and the residue partitioned between dichloromethane and water. The organic layer was washed with brine and dried over MgSO₄.


Concentration gave a solid, which was purified by chromatography on silica gel eluting with mixtures of dichloromethane and methanol.

LC retention time 4.50 minutes, $[M+H]^+$ 760.

- 5 b) The tris-Boc compound was treated with 1mL of 1/1 DCM/TFA and stirred at RT for 1h. The volatiles were removed *in vacuo* to give the desired mono-guanidine as the tris-trifluoroacetate.

LC retention time 2.90 minutes, $[M+H]^+$ 460.


10 Example 27



As example 26 using 4-*N,N'*-bis-*t*-butoxycarbonylguanidinoaminomethylaniline.

- 15 LC retention time 2.90 minutes, $[M+H]^+$ 460.


Example 28.



- 20 As example 26 using 4-aminomethylpyridine.

LC retention time 3.09 minutes, $[M+H]^+$ 404.

Example 29.




25

As example 26 using 2-(2-aminoethyl)pyridine.

LC retention time 3.19 minutes, $[M+H]^+$ 418.

Example 30.




5

As example 26 using 1-(3-aminopropyl)imidazole.

LC retention time 2.87 minutes, $[M+H]^+$ 421.

Example 31.




10

As example 26 using 1-(3-aminopropyl)morpholine.

LC retention time 2.86 minutes, $[M+H]^+$ 440.

15 Example 32.




As example 26 using 5-(2-aminoethyl)imidazole.

LC retention time 2.86 minutes, $[M+H]^+$ 407.

20


Example 32.



As example 26 using N,N-1,3-diaminopropane.

25 LC retention time 2.84 minutes, $[M+H]^+$ 398.


Example 33.



As example 26 using 4-aminomethylbenzenesulfonamide.

- 5 LC retention time 3.15 minutes, $[M+H]^+$ 482.


Example 34.



- 10 As example 26 using mono-*N,N'*-bis-Boc-piperazine carboxamidine.

LC retention time 2.92 minutes, $[M+H]^+$ 424.

Example 35.




15

As example 26 using mono-*N,N'*-bis-Boc-homopiperazine carboxamidine.

LC retention time 2.83 minutes, $[M+H]^+$ 438.

Example 36.




20

As example 26 using 5-aminomethyl-2-*N*-Boc-aminobenzimidazole.

LC retention time 2.89 minutes, $[M+H]^+$ 458.


Example 37.



As example 26 using 1-(3-aminopropyl)-2-methylpiperidine.

LC retention time 2.93 minutes, $[M+H]^+$ 452.

10 Example 38.




As example 26 using 1-(2-aminoethyl)pyrrolidine.

LC retention time 2.90 minutes, $[M+H]^+$ 410.

15


Example 39.



As example 26 using 4-(2-aminoethyl)benzene sulfonamide.

20 LC retention time 3.18 minutes, $[M+H]^+$ 496.2


Example 40.



25 As example 26 using *N,N*-dimethyl-1,2-diaminoethane.

LC retention time 2.89 minutes, $[M+H]^+$ 384.


Example 41.



As example 26 using tryptamine.


- 5 LC retention time 3.36 minutes, $[M+H]^+$ 456.

Example 42.




- 10 As example 26 using 1-(2-aminoethyl)-4-*N*-Boc-piperazine.
LC retention time 2.83 minutes, $[M+H]^+$ 425.

Example 43.




- 15 As example 26 using 1-(3-aminopropyl)-4-methylpiperazine.
LC retention time 2.86 minutes, $[M+H]^+$ 453.

Example 44.



- 20 As example 26 using 1-(3-aminopropyl)pyrrolidine.
LC retention time 2.87 minutes, $[M+H]^+$ 424.


Example 45.



As example 26 using 2-aminoethyl-1-ethylpyrrolidine.

- 5 LC retention time 2.95 minutes, $[M+H]^+$ 424.


Example 46.



- 10 As example 26 using *N,N*-diethyl-1,3-diaminopropane.

LC retention time 2.88 minutes, $[M+H]^+$ 426.

Example 47.




15

As example 26 using 4-aminomethyl-*N,N*-dimethylaniline.

LC retention time 3.34 minutes, $[M+H]^+$ 446.

Example 48.




20

As example 26 using *N,N*-dimethyl-2,2-dimethyl-1,3-diaminopropane.

LC retention time 2.93 minutes, $[M+H]^+$ 426.

Example 49.




As example 26 using 1-(3-aminopropyl)pyrrolidinone.

5 LC retention time 3.12 minutes, $[M+H]^+$ 438.

Example 50.

10




As example 26 using 4-N-Boc-piperazine glycinamide.

LC retention time 2.82 minutes, $[M+H]^+$ 439.

15


Example 51.



As example 26 using 4-aminomethyl-1-N-Boc-piperidine.

20 LC retention time 2.93 minutes, $[M+H]^+$ 410.


Example 52.



As example 26 using δ -*N*-Boc-D,L-lysine methyl ester.

LC retention time 2.94 minutes, $[M+H]^+$ 456.


Example 53.



As example 26 using 3-aminomethylpyridine.

LC retention time 2.93 minutes, $[M+H]^+$ 404.

10 Example 54.




As example 26 using 4-(2-aminoethyl)-*N*-Boc-piperidine.

LC retention time 2.87 minutes, $[M+H]^+$ 424.

15


Example 55.



As example 26 using 4-(2-aminoethyl)pyridine.

20 LC retention time 2.99 minutes, $[M+H]^+$ 418.

Example 56.




As example 26 using bis-(2-*N*-Boc-aminoethyl)amine.

LC retention time 0.62 minutes, $[M+H]^+$ 427.

Example 57

5




To a mixture of the aldehyde (1 eq) of example 10, dichlorobis(triphenylphosphine)-palladium(II) (10 mol%) and 2N Na_2CO_3 was added 1-*N*-*t*-butoxycarbonylindoline-5-boronic acid (2 eq) in dry, degassed DME and the mixture heated at 80°C for 1h. After cooling to RT the solvent was removed *in vacuo* and the residue partitioned between ethyl acetate and water. The organic layer was washed with brine and dried over MgSO_4 . Concentration gave a solid, which was purified by chromatography on silica gel eluting with mixtures of ethyl acetate and hexane.


15 LC retention time 5.23 minutes, $[M+H-\text{Boc}]^+$ 397.

Example 58

20



a)



25 The aldehyde (1 eq) from example 57 and 1-(3-aminopropyl)-2-methylpiperidine (1.6) were stirred at RT for 15min in 1,2-dichloroethane, then sodium triacetoxyborohydride (1.5) was added in one portion. After stirring for 16h at RT the mixture was concentrated


in vacuo and the residue partitioned between dichloromethane and water. The organic layer was washed with brine and dried over MgSO_4 . Concentration gave a solid, which was purified by chromatography on silica gel eluting with mixtures of dichloromethane and methanol.

5 LC retention time 3.47 minutes, $[\text{M}+\text{H}]^+$ 637.

b) The bis-Boc compound was treated with 1 mL of 1/1 DCM/TFA and stirred at RT for 10 min. The volatiles were removed *in vacuo* to give the desired compound as the tris-trifluoroacetate.

10 LC retention time 0.58 minutes, $[\text{M}+\text{H}]^+$ 437.

Example 59




15

As example 58 using *N,N*-diethyldiaminopropane.

LC retention time 0.52 and 0.84 minutes, $[\text{M}+\text{H}]^+$ 411.

Example 60

20




As example 58 using 1-(3-aminopropyl)pyrrolidinone.

LC retention time 0.56 and 2.60 minutes, $[\text{M}+\text{H}]^+$ 423.

25

Example 61




As example 58 using 1-(3-aminopropyl)imidazole.

LC retention time 0.54 and 0.78 minutes, $[M+H]^+$ 406.

5


Example 62



As example 58 using 4-(2-aminoethyl)pyridine.

LC retention time 0.57 minutes, $[M+H]^+$ 403.


15 Example 63




To a mixture of the aldehyde (1 eq) of example 10, dichlorobis(triphenylphosphine)-
20 palladium(II) (10 mol%) and 2N Na_2CO_3 was added 1-*N*-*t*-butoxycarbonyl-7-nitroindoline-5-boronic acid (2 eq) in dry, degassed DME and the mixture heated at 80°C for 1h. After cooling to RT the solvent was removed *in vacuo* and the residue partitioned between ethyl acetate and water. The organic layer was washed with brine and dried over MgSO_4 . Concentration gave a solid, which was purified by chromatography on silica gel
25 eluting with mixtures of ethyl acetate and hexane.

LC retention time 5.03 minutes, $[M+H-\text{Boc}]^+$ 442.

Example 64



5 a)




The aldehyde (1 eq) from example 63 and 1-(3-aminopropyl)-2-methylpiperidine (1.6 eq) were stirred at RT for 15min in 1,2-dichloroethane, then sodium triacetoxyborohydride (1.5 eq) was added in one portion. After stirring for 16h at RT the mixture was concentrated *in vacuo* and the residue partitioned between dichloromethane and water. The organic layer was washed with brine and dried over MgSO_4 . Concentration gave a solid, which was purified by chromatography on silica gel eluting with mixtures of dichloromethane and methanol.

LC retention time 4.10 minutes, $[\text{M}+\text{H}]^+$ 754.

b) The bis-Boc compound was treated with 1mL of 1/1 DCM/TFA and stirred at RT for 1h. The volatiles were removed *in vacuo* to give the desired compound as the trifluoroacetate.

LC retention time 2.84 minutes, $[\text{M}+\text{H}]^+$ 482.

Example 65




25

As example 64 using 1-(3-aminopropyl)imidazole.

LC retention time 2.82 minutes, $[M+H]^+$ 451.

Example 66

5




As example 64 using 4-(2-aminoethyl)-1-*N*-Boc-piperidine.

LC retention time 2.81 minutes, $[M+H]^+$ 454.

10


Example 67




- 15 To a mixture of the aldehyde (1 eq) of example 10, dichlorobis(triphenylphosphine)-palladium(II) (10 mol%) and 2N Na_2CO_3 was added 1-*N*-*t*-butoxycarbonyl-7-nitroindole-5-boronic acid in (2 eq) dry, degassed DME and the mixture heated at 80°C for 1h. After cooling to RT the solvent was removed *in vacuo* and the residue partitioned between ethyl acetate and water. The organic layer was washed with brine and dried over MgSO_4 .
- 20 Concentration gave a solid, which was purified by chromatography on silica gel eluting with mixtures of ethyl acetate and hexane.
- LC retention time 5.20 minutes, $[M+H-\text{Boc}]^+$ 440.

Example 68

25



a)



- 5 The aldehyde (1 eq) from example 67 and 1-(3-aminopropyl)-2-methylpiperidine (1.6 eq) were stirred at RT for 15min in 1,2-dichloroethane, then sodium triacetoxyborohydride (1.5 eq) was added in one portion. After stirring for 16h at RT the mixture was concentrated *in vacuo* and the residue partitioned between dichloromethane and water. The organic layer was washed with brine and dried over MgSO_4 . Concentration gave a
- 10 solid, which was purified by chromatography on silica gel eluting with mixtures of dichloromethane and methanol.


LC retention time 3.50 minutes, $[\text{M}+\text{H}]^+$ 680.

- b) The bis-Boc compound was treated with 1mL of 1/1 DCM/TFA and stirred at RT for 10
- 15 min. The volatiles were removed *in vacuo* to give the desired compound as the trifluoroacetate.

LC retention time 2.88 minutes, $[\text{M}+\text{H}]^+$ 480.

Example 69

20




As example 68 using 1-(3-aminopropyl)imidazole.

LC retention time 2.84 minutes. $[\text{M}+\text{H}]^+$ 449.

25

Example 70

44




As example 68 using 4-(2-aminoethyl)-1N-*t*-butoxycarbonylpiperidine.

LC retention time 2.85 minutes, $[M+H]^+$ 452.

5


Example 71



10 As example 68 using 4-(2-aminoethyl)pyridine.


LC retention time 2.95 minutes, $[M+H]^+$ 446.

Example 72



15

a)



20


To a mixture of the aldehyde (1 eq) of example 10, dichlorobis(triphenylphosphine)-palladium(II) (10 mol%) and 2N Na_2CO_3 was added phenylboronic acid (2 eq) in dry, degassed DME and the mixture heated at 80°C for 16h. After cooling to RT the solvent was removed *in vacuo* and the residue partitioned between ethyl acetate and water. The

organic layer was washed with brine and dried over MgSO_4 . Concentration gave a solid, which was purified by chromatography on silica gel eluting with mixtures of ethyl acetate and hexane.

LC retention time 4.81 minutes, $[\text{M}+\text{Na}]^+$ 378.

5

b)



- 10 The aldehyde (1 eq) and mono-*N,N'*-bis-*t*-butoxycarbonylpiperazinecarboxamidine (1.6) were stirred at RT for 15min in 1,2-dichloroethane, then sodium triacetoxyborohydride (1.5) was added in one portion. After stirring for 16h at RT the mixture was concentrated *in vacuo* and the residue partitioned between dichloromethane and water. The organic layer was washed with brine and dried over MgSO_4 . Concentration gave a solid, which
- 15 was purified by chromatography on silica gel eluting with mixtures of hexane and ethyl acetate.


LC retention time 4.03 minutes, $[\text{M}+\text{H}]^+$ 668.

- c) The tris-Boc compound was treated with 1mL of 1/1 DCM/TFA and stirred at RT for
- 20 1h. The volatiles were removed *in vacuo* to give the desired compound as the tris-trifluoroacetate.

LC retention time 0.52 and 2.57 minutes, $[\text{M}+\text{H}]^+$ 368.

Example 73

25




As example 72 using *p*-tolylboronic acid.

LC retention time 2.71 minutes, $[M+H]^+$ 382.

Example 74

5




As example 72 using 4-methoxyphenylboronic acid.

LC retention time 2.64 minutes, $[M+H]^+$ 398.

10


Example 75



15 As example 72 using 4-fluorophenylboronic acid.

LC retention time 2.63 minutes, $[M+H]^+$ 386.

Example 76




20

As example 72 using 3-trifluoromethylphenylboronic acid.

LC retention time 2.86 minutes. $[M+H]^+$ 436.


Example 77




As example 72 using 3-thienylboronic acid.

LC retention time 0.51 and 2.50 minutes, $[M+H]^+$ 374.

10 Example 78



a)




To a mixture of the aldehyde (1 eq) of example 10, dichlorobis(triphenylphosphine)-palladium(II) (10 mol%) and 2N Na₂CO₃ was added phenylboronic acid (2 eq) in dry, degassed DME and the mixture heated at 80°C for 16h. After cooling to RT the solvent was removed *in vacuo* and the residue partitioned between ethyl acetate and water. The organic layer was washed with brine and dried over MgSO₄. Concentration gave a solid, which was purified by chromatography on silica gel eluting with mixtures of ethyl acetate and hexane.

LC retention time 4.81 minutes. $[M+Na]^+$ 378.

25

b)



- 5 The aldehyde (1 eq) and mono-*N,N'*-bis-*t*-butoxycarbonylhomopiperazinecarboxamidine (1.6 eq) were stirred at RT for 15min in 1,2-dichloroethane, then sodium triacetoxyborohydride (1.5 eq) was added in one portion. After stirring for 16h at RT the mixture was concentrated *in vacuo* and the residue partitioned between dichloromethane and water. The organic layer was washed with brine and dried over MgSO_4 .
- 10 Concentration gave a solid, which was purified by chromatography on silica gel eluting with mixtures of hexane and ethyl acetate.


LC retention time 4.03 minutes, $[\text{M}+\text{H}]^+$ 668.

- c) The tris-Boc compound was treated with 1mL of 1/1 DCM/TFA and stirred at RT for 15 1h. The volatiles were removed *in vacuo* to give the desired compound as the tris-trifluoroacetate.

LC retention time 0.50 and 2.46 minutes, $[\text{M}+\text{H}]^+$ 382.

Example 79

20




As example 78 using *p*-tolylboronic acid.

LC retention time 2.65 minutes. $[\text{M}+\text{H}]^+$ 396.


25

Example 80



- 5 As example 78 using 4-methoxyphenylboronic acid.
LC retention time 0.53 and 2.59 minutes, $[M+H]^+$ 412.


Example 81



10


- As example 78 using 4-fluorophenylboronic acid.
LC retention time 0.54 and 2.56 minutes, $[M+H]^+$ 400.

15 Example 82




20

Example 83



As example 78 using 3-thienylboronic acid.

LC retention time 0.50 and 2.10 minutes, $[M+H]^+$ 388.



5


Reagents: (i) R^2NH_2 , DCE, sodium triacetoxyborohydride; (ii) Di-*t*-butyl dicarbonate, DCM, (i-Pr) $_2$ NEt; (iii) Bis(pinacolato)diboron, DMSO, KOAc, $PdCl_2(dppf)_2$; (iv) Arylboronic acid, DMF, K_3PO_4 , $PdCl_2(dppf)_2$; (v) TFA/DCM, 1/1.

Scheme 4

10


The following examples were synthesized following the procedure outlined in Scheme 4.

Example 84



15


a)




The aldehyde (1eq.) from example 10 and 4-(aminomethyl)-1-Boc-piperidine (1.6eq.) were stirred at RT for 15min in 1,2-dichloroethane. then sodium triacetoxymethylborohydride (1.5eq.) was added in one portion. After stirring for 16 hours at RT the mixture was concentrated *in vacuo* and the residue partitioned between EtOAc and water. The organic layer was washed with brine and dried over MgSO_4 . Concentration gave a solid, which was dissolved in DCM and diisopropylethylamine (1eq.) and di-*t*-butyl dicarbonate (3eq.) added. The reaction was stirred at RT for 1h, diluted with water and the organic layer separated and washed with brine. Drying (MgSO_4) and concentration *in vacuo* gave a solid, which was purified by chromatography on silica gel eluting with ethyl acetate/hexane mixtures. LC retention time 5.41minutes, $[\text{M-Boc}+\text{H}]^+$ 556.

b) The bromophenyl compound (1eq.), bis(pinacolato)diboron (1.1eq.) and potassium acetate (3eq.) in DMSO were treated with bis(diphenylphosphino)ferrocene palladium dichloride (10mol %) and heated at 80°C for 3h. After this time, the solvents were partitioned between water and diethylether. the organic layer dried with MgSO_4 and finally concentrated. The oil produced was purified using chromatography on silica gel using ethyl acetate/hexanes. LC retention time 5.45minutes, $[\text{M-Boc}+\text{H}]^+$ 604.

Example 85



25 a)




The boronate ester (1eq.) from example 84 was treated with 5-bromo-1-N-Boc-indole (1.5eq.), potassium phosphate (3eq.) and bis(diphenylphosphino)ferrocene palladium dichloride (10 mol%) in DMF at 60°C for 4h. The mixture was filtered through celite and concentrated *in vacuo*. The residue was partitioned between EtOAc and water, the organic layer was washed with brine and dried (MgSO₄). Concentration gave a residue which was purified by column chromatography on silica eluting with ethyl acetate/hexane mixtures.

LC retention time 5.87 minutes, [M-Boc+H]⁺ 694.

- 10 b) The fully protected compound was treated with 1mL of 1/1 DCM/TFA and stirred at RT for 10min. The volatiles were removed *in vacuo* to give the desired compound as the tris-trifluoroacetate.

LC retention time 2.68 minutes, [M+H]⁺ 392.

15 Example 86




As example 85 using 5-bromobenzofuran-2-methylamide.

LC retention time 2.67 minutes, [M+H]⁺ 451.

20


Example 87



As example 85 using 2-bromopyridine.

- 25 LC retention time 0.45 minutes, [M+H]⁺ 355.


Example 88



As example 85 using 5-bromoindan-2-one.


5 LC retention time 3.09 minutes, $[M+H]^+$ 408.

Example 89




10 As example 85 using 3-bromo(4'-fluoro)diphenylether.
LC retention time 3.24 minutes, $[M+H]^+$ 464.

Example 90



15
As example 85 using 5-bromochromanone.
LC retention time 3.10 minutes, $[M+H]^+$ 422.


Example 91



20
As example 85 using 3-bromoquinoline.
LC retention time 3.09 minutes, $[M+H]^+$ 405.

25 Example 92

54




As example 85 using bromobenzene.

LC retention time 0.49 minutes, $[M+H]^+$ 354.

5


Example 93



As example 85 using 6-bromoquinoline.

10 LC retention time 0.47 minutes, $[M+H]^+$ 419.


Example 94



15 As example 85 using 5-bromo-2,4-dimethoxypyrimidine.

LC retention time 3.07 minutes, $[M+H]^+$ 416.

Example 95




20

As example 85 using 5-bromoindole-2-methylamide.

LC retention time 3.22 minutes. $[M+H]^+$ 450.

25


Example 96




As example 85 using 5-bromoindole-2-benzylamide.

5 LC retention time 1.11 minutes, $[M+H]^+$ 526.

Example 97



10 a)




The aldehyde (1eq.) from example 10 and 4-(2-aminomethyl)-1-Boc-piperazine (1.5eq.) were stirred at RT for 15min in 1,2-dichloroethane, then sodium triacetoxyborohydride (1.5eq.) was added in one portion. After stirring 16 hours at RT the mixture was concentrated *in vacuo* and the residue partitioned between EtOAc and water. The organic layer was washed with brine and dried over $MgSO_4$. Concentration gave a solid which was dissolved in DCM, and diisopropylethylamine (5eq) and di-*t*-butyl dicarbonate (2eq.) added. The reaction was stirred at RT for 1h. diluted with water and the organic layer separated and washed with brine. Drying ($MgSO_4$) and concentration *in vacuo* gave a solid which was purified by chromatography on silica gel eluting with ethyl acetate/hexane mixtures.

LC retention time 4.25 minutes. $[M+H]^+$ 673.

b) The bromophenyl compound (1eq.), bis(pinacolato)diboron (1.1eq.) and potassium acetate (3eq.) in DMSO were treated with bis(diphenylphosphino)ferrocene palladium dichloride (10 mol%) and heated at 80°C for 3h. After this time, the solvents were partitioned between water and diethylether, the organic layer dried with MgSO₄ and finally concentrated. The oil produced was purified using chromatography on silica gel using ethyl acetate/hexanes.


LC retention time 4.27 minutes, [M+H]⁺ 719.

Example 98



10

a)



15

The boronate ester (1eq.) from example 97 was treated with 5-bromo-1-Boc-indole (1.5eq.), potassium phosphate (3eq.) and with bis(diphenylphosphino)ferrocene palladium dichloride (10 mol%) in DMF at 60°C for h. The mixture was filtered through celite and concentrated *in vacuo*. The residue was partitioned between between EtOAc and water, the organic layer was washed with brine and dried (MgSO₄). Concentration gave a residue which was purified by column chromatography on silica gel eluting with ethyl acetate/hexane mixtures.

20

LC retention time 4.79 minutes. [M+H]⁺ 808.


25

b) The fully protected compound was treated with 1mL of 1/1 DCM/TFA and stirred at RT for 10min. The volatiles were removed *in vacuo* to give the desired compound as the tris-trifluoroacetate.

LC retention time 2.70 minutes, $[M+H]^+$ 408.

Example 99


5



As example 98 using 5-bromo-2,4-dimethoxypyrimidine.

LC retention time 1.03 minutes, $[M+H]^+$ 431.

10 Example 100




As example 98 using 1,2,3,4-tetrahydro-9-Boc-6-bromocarbazole.

LC retention time 3.21 minutes, $[M+H]^+$ 462.

15

58




Reagents: (i) 3-bromo-1-phthalamidopropane, Cs_2CO_3 , DMF; (ii) 2-Benzo[b]thiopheneboronic acid, DME, K_2CO_3 ; (iii) (a) (*N,N'*-bis-butoxycarbonyl) $\text{NH}(\text{CH}_2)_4\text{NH}_2$, DCE, sodium triacetoxymethylborohydride, (b) di-*t*-butyl dicarbonate, DCM, (i-Pr) $_2\text{NEt}$; (iv) hydrazine hydrate, ; (v) RSO_2Cl , Et_3N , DCM; (vi) RCOCl , Et_3N , CH_3CN ; (vii) RCl (R =heterocycle), (i-Pr) $_2\text{NEt}$, ; (viii) TFA/DCM, 1/1.


Scheme 5

10 The following examples were synthesized following the procedure outlined in Scheme 5.

Example 101.



a)




5-Bromosalicylaldehyde (1 eq.) and cesium carbonate (2 eq.) were stirred in DMF and 3-bromo-1-phthalamidopropane (1.2 eq.) was added. After stirring for 18 hours at room temperature the reaction was concentrated *in vacuo*. The residue was partitioned between EtOAc and water and the organic layer washed with brine. Drying (MgSO_4) and concentration gave a residue which was purified by column chromatography on silica eluting with EtOAc/hexane to yield the protected amine.

LC retention time 4.69 minutes, $[\text{M}]^+$ 388.

10

b)




To the aldehyde (1 eq.), 2N K_2CO_3 and benzo[b]thiophene-2-boronic acid (2 eq.) in dry, degassed DME was added dichlorobis(triphenylphosphine)palladium (0.1 eq.), and the mixture heated at 80°C for 18 h. After this time the reaction mixture was filtered, concentrated *in vacuo*, re-dissolved in ethyl acetate and washed with water and saturated sodium chloride solution. The organic extract was dried (MgSO_4), filtered and concentrated *in vacuo*.

LC retention time 5.24 minutes, $[\text{M}+\text{H}]^+$ 442.

20


c)



The aldehyde (1 eq.) and 1-amino-4-*N,N'*-bis-Boc-guanidinobutane (1.5 eq.) were stirred at RT for 15min in 1,2-dichloroethane, then sodium triacetoxyborohydride (1.5 eq.) was added in one portion. After stirring for 18 hours at RT the mixture was concentrated *in vacuo* and the residue partitioned between EtOAc and water. The organic layer was washed with brine and dried over MgSO_4 . Concentration gave a solid which was dissolved in DCM and treated with diisopropylethylamine (2 eq.) and di-*t*-butyl dicarbonate (2 eq.). After stirring for 4h at RT the reaction was diluted with water, the organic layer was washed with brine, and dried over MgSO_4 . Concentration gave a solid which was purified by chromatography on silica gel eluting with 4:1 hexane:ethyl acetate, then 2:1 hexane:ethyl acetate

LC retention time 5.95 minutes, $[\text{M}+\text{H}]^+$ 856.4

d)



The *N*-protected phthalimide (1 eq.) was treated with hydrazine hydrate (15 eq.) in EtOH for 18 h then concentrated *in vacuo* keeping the water bath below 40°C. The residue was dissolved in EtOAc and the organic layer washed exhaustively with water and finally with brine. The organic fraction was dried (MgSO_4), concentrated *in vacuo* and purified by column chromatography on silica, eluting with 75%EtOAc/hexane, then 5%MeOH/DCM.


20

e) The amine (1 eq.) and triethylamine (5 eq.) in acetonitrile was treated with methanesulfonyl chloride (3 eq.). After 2h at RT the reaction was diluted with water and the organic layer separated, washed with brine and dried (MgSO_4). Concentration gave an oil which was purified by column chromatography on silica, eluting with 1:1 EtOAc/hexane to yield the desired methane sulfonate.

The bis-Boc compound was treated with 1 mL of 1/1 DCM/TFA and stirred at RT for 1h. The volatiles were removed *in vacuo* to give the desired mono-guanidine as the bis-trifluoroacetate.

LC retention time 3.19 minutes. $[\text{M}-\text{H}]^-$ 504.


Example 102.



As example 101 using benzenesulfonyl chloride.

5 LC retention time 3.35 minutes, $[M+H]^+$ 566.

Example 103.




10 As example 101 using acetyl chloride.

LC retention time 3.12 minutes, $[M+H]^+$ 468.

15


Example 104.



As example 101 using 2-chloro-4,6-dimethoxytriazine.


20 LC retention time 3.24 minutes. $[M+H]^+$ 565.

Example 105.



As example 101 using 2-chloro-4-trifluoromethylpyrimidine.

5 LC retention time 3.45 minutes, $[M+H]^+$ 518.




Reagents: (i) Bis(pinacolato)diboron, KOAc, DMSO, $\text{PdCl}_2(\text{dppf})_2$; (ii) Arylboronic acid,

10 DMF, K_3PO_4 , $\text{PdCl}_2(\text{dppf})_2$; (iii) $(\text{N},\text{N}'\text{-bis-}t\text{-butoxycarbonyl})\text{NH}(\text{CH}_2)_3\text{CH}_2\text{NH}_2$, DCE, sodium triacetoxyborohydride; (iv) TFA/DCM, 1/1.

Scheme 6

15 The following examples were synthesized following the procedure outlined in Scheme 6.


Example 106




The aldehyde (1eq.) from example 10, bis(pinacolato)diboron (1.1eq.) and potassium acetate (3eq.) in DMSO were treated with bis(diphenylphosphino)ferrocene palladium dichloride (10 mol%) and heated at 80°C for 3h. After this time, the solvents were partitioned between water and diethylether, the organic layer dried with MgSO₄ and finally concentrated. The oil produced was purified using chromatography on silica gel using ethyl acetate/hexanes.

LC retention time 4.81 minutes, [M-Boc+H]⁺ 306.

Example 107



a)




15 The boronate ester (1eq.) from example 106 was treated with 5-bromobenzofuran-2-methylamide (1.5eq.), potassium phosphate (3eq.) and bis(diphenylphosphino)ferrocene palladium dichloride (10 mol%) in DMF at 60°C for 3h. The mixture was filtered through celite and concentrated *in vacuo*. The residue was partitioned between EtOAc and water, the organic layer was washed with brine and dried (MgSO₄). Concentration gave a

20 residue which was purified by column chromatography on silica eluting with ethyl acetate/hexane mixtures.

LC retention time 4.89 minutes, [M-Boc+H]⁺ 429.

b)




The aldehyde (1eq.) and 1-amino-4-(N,N'-bis-Boc-guanidino)butane (1.5eq.) were stirred at RT for 15min in dichloroethane, then sodium triacetoxyborohydride (1.5eq.) was added in one portion. After stirring 16 hours at RT the mixture was concentrated *in vacuo* and the residue partitioned between EtOAc and water. The organic layer was washed with brine and dried over MgSO₄. Concentration gave a solid which was purified by chromatography on silica gel eluting with ethyl acetate/hexane mixtures.

LC retention time 4.40 minutes, [M-Boc+H]⁺ 667.

c) The tris-Boc compound was treated with 1mL of 1/1 DCM/TFA and stirred at RT for 1h. The volatiles were removed *in vacuo* to give the desired compound as the tris-trifluoroacetate.

LC retention time 3.17 minutes, [M+H]⁺ 467.

15 Example 108




As example 107 using 5-bromobenzofuran-2-benzylamide.

LC retention time 3.19 minutes, [M+H]⁺ 543.

20


Example 109



As example 107 using 1-bromo-2,3-dimethoxybenzene.

25 LC retention time 3.10 minutes, [M+H]⁺ 430.


Example 110



As example 107 using 2-bromothiazole.

5 LC retention time 1.06 minutes, $[M+H]^+$ 377.

Example 111




10

As example 107 using 3-bromoquinoline.

LC retention time 3.13 minutes, $[M+H]^+$ 421.

Example 112




15

As example 107 using 5-bromo-1-butoxycarbonylindole.


LC retention time 3.13 minutes, $[M+H]^+$ 409.

20 Example 113



As example 107 using 5-bromo-1-methylindole.

LC retention time 2.77 minutes, $[M+H]^+$ 423.




Reagents: (i) BnBr, Cs₂CO₃, DMF ; (ii) ArB(OH)₂, DME, K₂CO₃, PdCl₂(PPh₃)₂ ; (iii) BCl₃·SMe₂, DCM; (iv) R₁R₂NH, MeOH-CH₂Cl₂ (1:1 v/v), Amberlyst A-26 borohydride resin; (v) Boc₂O, DIPEA, MeCN; (vi) EtOH, Cs₂CO₃, then R³X, DMF.

Scheme 7


The following examples were synthesised following the procedure outlined in scheme 7.

10

Example 114.



a)




15

To 5-bromosalicylaldehyde (1 eq.) and Cs₂CO₃ (2 eq.) in DMF was added benzyl bromide (1.1 eq.), and the solution stirred under nitrogen for 24 hours. The mixture was poured into water and extracted with ethyl acetate. The organic fraction was washed with water and saturated sodium chloride solution, dried (MgSO₄), filtered and concentrated *in vacuo* to give the benzyl ether as a white solid.

20

b)




To the aryl bromide (1 eq.), 2N K_2CO_3 and benzo[b]thiophene-2-boronic acid (2 eq.) in dry, degassed DME was added dichlorobis(triphenylphosphine)palladium(II) (0.1 eq.) and the mixture heated at 80°C for 45 hours. After this time the reaction mixture was filtered, concentrated *in vacuo*, re-dissolved in ethyl acetate and washed with water and saturated sodium chloride solution. The organic extract was dried ($MgSO_4$), filtered and concentrated *in vacuo*.


LC retention time 5.47 minutes, $[M+H]^+$ 345.

c) The bisaryl (example 114b, 1 eq.) and boron trichloride-methyl sulfide complex (2 eq.) were stirred in dichloromethane for one hour. Saturated sodium bicarbonate was added and the organics collected. After passing through a short pad of silica and concentration under reduced pressure, the product was suspended in methanol-dichloromethane (1:1 v/v) and shaken with 4-(aminomethyl)piperidine (1 eq.). After one hour, borohydride (solid supported upon Amberlyst A-26 resin, 1.1 eq.) was added and shaking continued for 18 hours. After filtration, the solution was concentrated under reduced pressure, re-dissolved in acetonitrile and stirred for 18 hours with di-*t*-butyl dicarbonate (2 eq.) and diisopropylethylamine (2 eq.). The resultant mixture was poured into water and extracted with ethyl acetate. After concentration *in vacuo*, the compound was purified by chromatography on silica gel, eluting with hexanes and ethyl acetate (4:1 v/v).
LC retention time 5.68 minutes.

25 Example 115



a)




The bisaryl (example 114, 1 eq.) and Cs_2CO_3 (2 eq.) in ethanol were heated to 80°C for one hour, and then concentrated to dryness. After re-suspension in DMF and warming to 45°C under nitrogen, 3-(3-pyridyl)-1-bromopropane (1.5 eq.) was added and the suspension stirred for 18 hours. After concentration *in vacuo*, the compound was purified by chromatography on silica gel, eluting with hexane:ethyl acetate (2:1 v/v), then dichloromethane:methanol (9:1 v/v).

10

b) The thus purified fully protected compound was treated with 1 mL of 1/1 TFA/DCM and stirred at RT for 30 minutes. The volatiles were removed under reduced pressure to give the desired diamine as the tris-trifluoroacetate.

15 LC retention time 3.02 minutes, $[\text{M}+\text{H}]^+$ 472.


Example 116.



20 As example 115 using 1-chloro-3-(N,N-dimethylamino)propane.

LC retention time 3.02 minutes, $[\text{M}+\text{H}]^+$ 438.


Example 117.



As example 115 using 2-(2-chloroethyl)-1-methylpyrrolidine.


- 5 LC retention time 3.00 minutes, $[M+H]^+$ 464.

Example 118.



- 10 As example 115 using 1-chloro-3-(N,N-dimethylamino)-2-methylpropane.
LC retention time 3.16 minutes, $[M+H]^+$ 452.

Example 119.




15

As example 115 using 4-(2-chloroethyl)morpholine.

LC retention time 3.17 minutes, $[M+H]^+$ 466.

20


Example 120.



As example 115 using N-(2-chloroethyl)pyrrolidine.

5 LC retention time 3.16 minutes, $[M+H]^+$ 450.


Example 121.



10 As example 115 using 2-bromoethyl-N,N-diethylamine.

LC retention time 3.20 minutes, $[M+H]^+$ 452.

Example 122.




15

As example 115 using N-Boc¹-3-(bromomethyl)piperidine.

LC retention time 3.24 minutes. $[M+H]^+$ 450.

20


Example 123.



As example 115 using *N*-Boc^t-4-(2-bromoethyl)piperidine.

5 LC retention time 3.24 minutes, $[M+H]^+$ 464.

Example 124




10


As example 114 using *N*-(2-aminoethyl)piperidine.

LC retention time 4.75 minutes, $[M+H]^+$ 668.

15 Example 125.



a)




The bisaryl (example 124, 1 eq.) and Cs_2CO_3 (2 eq.) in ethanol were heated to 80°C for one hour, and then concentrated to dryness. After re-suspension in DMF and warming to 45°C under nitrogen, 3-(3-pyridyl)-1-bromopropane (1.5 eq.) was added and the suspension stirred for 18 hours. After concentration *in vacuo*, the compound was purified by chromatography on silica gel, eluting with hexane:ethyl acetate (2:1 v/v), then dichloromethane:methanol (9:1 v/v).

b) The thus purified fully protected compound was treated with 1 mL of 1/1 TFA/DCM and stirred at RT for 30 minutes. The volatiles were removed under reduced pressure to give the desired triamine as the tetrakis-trifluoroacetate.


LC retention time 3.09 minutes. $[\text{M}+\text{H}]^+$ 487.

Example 126.



As example 125 using 1-chloro-3-(N,N-dimethylamino)propane
LC retention time 2.91 minutes, $[\text{M}+\text{H}]^+$ 453.


20 Example 127.



As example 125 using 2-(2-chloroethyl)-1-methylpyrrolidine.
LC retention time 2.97 minutes. $[\text{M}+\text{H}]^+$ 479.

25

Example 128.




As example 125 using 2-bromoethyl-N,N-diethylamine.

LC retention time 3.21 minutes, $[M+H]^+$ 467.

5


Example 129.



As example 125 using 1-chloro-3-(N,N-dimethylamino)-2-methylpropane.

10 LC retention time 3.21 minutes, $[M+H]^+$ 467.


Example 130.



15 As example 125 using 2-picolyl chloride.

LC retention time 3.30 minutes, $[M+H]^+$ 459.

Example 131.




20

As example 125 using 3-picolyl chloride.

LC retention time 3.27 minutes, $[M+H]^+$ 459.

Example 132.




5

As example 125 using 4-picolyl chloride.

LC retention time 3.26 minutes, $[M+H]^+$ 459.

Example 133.




10

As example 125 using 4-(2-chloroethyl)morpholine.

LC retention time 3.18 minutes, $[M+H]^+$ 481.


15 Example 134.



As example 125 using 2-amino-6-(bromoethanoylamino)pyridine.

LC retention time 3.34 minutes, $[M+H]^+$ 517.

20




5


Scheme 8

The following examples were synthesised following the procedure outlined in scheme 8.

10 Example 135.



a)



15


To 5-bromosalicylaldehyde (1 eq.) and Cs_2CO_3 (2 eq.) in DMF was added Boc-3-bromopropylamine (1.2 eq.), and the solution stirred under nitrogen for 24 hours. The

mixture was poured into water and extracted with ethyl acetate. The organic fraction was washed with water and saturated sodium chloride solution, dried and concentrated *in vacuo* to give the ether as a white solid.

LC retention time 4.68 minutes, $[M+H]^+$ 380.

5

b)




To the aryl bromide (1 eq.), 2N K_2CO_3 and benzo[b]thiophene-2-boronic acid (2 eq.) in dry, degassed DME was added dichlorobis(triphenylphosphine)palladium(II) (0.1 eq.) and the mixture heated at 80°C for 45 hours. After this time the reaction mixture was filtered, concentrated *in vacuo*, re-dissolved in ethyl acetate and washed with water and saturated sodium chloride solution. The organic extract was dried ($MgSO_4$), filtered and concentrated *in vacuo*.

15 LC retention time 5.19 minutes, $[M-Boc^t+H]^+$ 298.


c) The aldehyde (1 eq.), hydroxylamine hydrochloride (1.5 eq.) and triethylamine (2 eq.) were stirred for 18 hours in 1,2-dichloroethane. The mixture was then poured into dichloromethane and washed with brine. The organic fraction was dried, filtered and concentrated under reduced pressure. The resultant compound was re-dissolved in methanol, a few drops of chloroform were added, and the mixture stirred under a hydrogen atmosphere for 18 hours over 10% palladium on carbon (1 eq. by weight). After filtration through a short pad of celite, the volatiles were removed under reduced pressure to give the amine as the hydrochloride salt.

25 LC retention time 3.75 minutes, $[M+H]^+$ 413.

Example 136.



a)




- 5 The amine from example 135, Boc-3-aminopropanoic acid (1 eq.), hydroxybenzotriazole (1.1 eq.) and triethylamine (5 eq.) were stirred together in DMF for 18 hours. The reaction mixture was concentrated *in vacuo* and partitioned between EtOAc and water and the organic layer dried (MgSO₄) and concentrated in *vacuo*. The residue was purified by column chromatography on silica eluting with 75%EtOAc/hexane.
- 10 LC retention time 5.05 minutes, [M+H]⁺ 570.

b) The thus purified fully protected compound was treated with 1 mL of 1/1 TFA/DCM and stirred at RT for 30 minutes. The volatiles were removed under reduced pressure to give the desired diamine as the bistrifluoroacetate.

15


Example 137.



As for example 136, using N-Boc-4-aminobutanoic acid.

20

78




- Reagents: (i) (a) TFA/DCM, (b) 2-Nitro-phenylsulfonyl chloride, TEA. DCM; (ii) RX.
 5 Cs₂CO₃, DMF; (iii) PhSH, DBU, CH₃CN; (iv) Boc₂O, DIPEA, CH₃CN; (v) 1-amino-4-(N,N'-bis-Bocguanidino)butane, NaB(OAc)₃H, 1,2-DCE; (vi) TFA/DCM, 1/1.

Scheme 9

- 10 The following examples were synthesised following the procedure outlined in scheme 9.

Example 138.




15

The aldehyde from example 25 (1 eq.) was dissolved in 1 mL of 1/1 TFA/DCM and stirred at RT for 30 minutes. The volatiles were removed under reduced pressure to give the amine as the trifluoroacetate salt. This salt was re-dissolved in dichloromethane and


stirred with 2-nitrophenylsulfonyl chloride (2 eq.) and triethylamine (2 eq.) for 2 hours. The mixture was poured into water and extracted with ethyl acetate. The organic fraction was washed with water and saturated sodium chloride solution, dried (MgSO_4) and concentrated under reduced pressure to give the sulfonamide as a white solid.

5

Example 139




a)



- 10 To the aldehyde from example 138 (1 eq.), and Cs_2CO_3 (2 eq.) in DMF was added methyl iodide (2 eq.), and the solution stirred under nitrogen for 24 hours. The mixture was poured into water and extracted with ethyl acetate. The organic fraction was washed with water and saturated sodium chloride solution, dried and concentrated under reduced pressure.

15


b)



- The biaryl (1 eq.), DBU (5 eq.) and thiophenol (4 eq.) were stirred together in DMF for 2h
20 hours. The mixture was concentrated *in vacuo* and partitioned between EtOAc and water. the organic layer dried (MgSO_4) and concentrated *in vacuo*. The compound was then re-dissolved in acetonitrile and stirred for 18 hours with di-*t*-butyl dicarbonate (2 eq.) and di-isopropylethylamine (2 eq.). The resultant mixture was poured into water and extracted

with ethyl acetate. The organic fraction was dried (MgSO_4), filtered and concentrated *in vacuo* to give the desired compound.

c)




The aldehyde (1 eq.) and 1-amino-4-(N,N'-bis-Boc-guanidino)butane (1.5 eq.) were stirred at RT for 15min in dichloroethane, then sodium triacetoxyborohydride (1.5 eq.) was added. After stirring for 18 hours at RT the mixture was concentrated *in vacuo* and the residue partitioned between EtOAc and water. The organic layer was dried, filtered and concentrated *in vacuo* to give the crude product. This was purified by column chromatography (silica, eluting with dichloromethane, then dichloromethane: methanol 95:5 v/v).

d) The thus purified fully protected compound was treated with 1 mL of 1/1 TFA/DCM and stirred at RT for 30 minutes. The volatiles were removed *in vacuo* to give the desired guanidine as the tris-trifluoroacetate.

LC retention time 2.88 minutes. $[\text{M}+\text{H}]^+ 440$.


20 Example 140.



As for example 139, using benzyl bromide.

25 LC retention time 3.054 minutes. $[\text{M}+\text{H}]^+ 516.25$.


Example 141



- 5 To a mixture of the aldehyde (1 eq) of example 10, dichlorobis(triphenylphosphine)-palladium(II) (10 mol%) and 2N Na₂CO₃ was added 5-methoxyindole-2-boronic acid (2 eq) in dry, degassed DME and the mixture heated at 80°C for 1h. After cooling to RT the solvent was removed *in vacuo* and the residue partitioned between ethyl acetate and water. The organic layer was washed with brine and dried over MgSO₄. Concentration gave a
- 10 solid, which was purified by chromatography on silica gel eluting with mixtures of ethyl acetate and hexane.
- LC retention time 5.19 minutes, [M+H-Boc]⁺ 425.


Example 142

15



a)

20



- The aldehyde (1 eq) from example 141 and N,N-Dimethylethylenediamine (1.6 eq) were stirred at RT for 15min in 1,2-dichloroethane, then sodium triacetoxyborohydride (1.5) was added in one portion. After stirring for 16h at RT the mixture was concentrated *in vacuo*
- 25 and the residue partitioned between dichloromethane and water. The organic layer was


washed with brine and dried over MgSO_4 . Concentration gave a solid, which was purified by chromatography on silica gel eluting with mixtures of dichloromethane and methanol.

LC retention time 3.65 minutes, $[\text{M}+\text{H}]^+$ 567.

- 5 b) The bis-Boc compound was treated with 1mL of 1/1 DCM/TFA and stirred at RT for 10 min. The volatiles were removed *in vacuo* to give the desired compound as the tris-trifluoroacetate.

LC retention time 2.81 minutes, $[\text{M}+\text{H}]^+$ 397.


10 Example 143



As example 142 using N,N-Diethylpropylenediamine.

- 15 LC retention time 2.80 minutes, $[\text{M}+\text{H}]^+$ 439.

Example 144




20

As example 142 using 1-(3-Aminopropyl)-2-pipecoline.

LC retention time 2.88 minutes. $[\text{M}+\text{H}]^+$ 465.


25

Example 145




- 5 As example 142 using 1-(2-Aminoethyl)piperazine.
LC retention time 2.82 minutes, $[M+H]^+$ 438.

Example 146



- As example 142 using 1-(3-Aminopropyl)imidazole.
LC retention time 2.78 minutes, $[M+H]^+$ 434.


15 Example 147




- To a mixture of the aldehyde (1 eq) of example 10, dichlorobis(triphenylphosphine)-
20 palladium(II) (10 mol%) and 2N Na_2CO_3 was added 3-methylindole-2-boronic acid (2 eq)
in dry, degassed DME and the mixture heated at 80°C for 1h. After cooling to RT the
solvent was removed *in vacuo* and the residue partitioned between ethyl acetate and water.
The organic layer was washed with brine and dried over MgSO_4 . Concentration gave a
solid, which was purified by chromatography on silica gel eluting with mixtures of ethyl
25 acetate and hexane.

LC retention time 5.33 minutes, $[M+H-Boc]^+$ 409.

Example 148



a)



10

The aldehyde (1 eq) from example 147 and 1-(3-Aminopropyl)-2-pipecoline (1.6 eq) were stirred at RT for 15min in 1,2-dichloroethane, then sodium triacetoxyborohydride (1.5) was added in one portion. After stirring for 16h at RT the mixture was concentrated *in vacuo* and the residue partitioned between dichloromethane and water. The organic layer was washed with brine and dried over $MgSO_4$. Concentration gave a solid, which was purified by chromatography on silica gel eluting with mixtures of dichloromethane and methanol.

15

LC retention time 3.65 minutes, $[M+H]^+$ 649.


b) The bis-Boc compound was treated with 1mL of 1/1 DCM/TFA and stirred at RT for 10 min. The volatiles were removed *in vacuo* to give the desired compound as the trifluoroacetate.

20

LC retention time 2.94 minutes. $[M+H]^+$ 449.

Example 149

25




As example 148 using 1-(2-Aminoethyl)piperazine.

LC retention time 2.88 minutes, $[M+H]^+$ 422.

Example 150

5




As example 148 using 1-(2-Aminoethyl)piperidine.

LC retention time 2.90 minutes, $[M+H]^+$ 421.

10

Example 151




15 As example 148 using 4-(Aminomethyl)piperidine.

LC retention time 3.58 minutes, $[M+H]^+$ 406.

Example 152

20




To a mixture of the aldehyde (1 eq) of example 10, dichlorobis(triphenylphosphine)-palladium(II) (10 mol%) and 2N Na_2CO_3 was added 4-methylindole-2-boronic acid (2 eq) in dry, degassed DME and the mixture heated at 80°C for 1h. After cooling to RT the

solvent was removed *in vacuo* and the residue partitioned between ethyl acetate and water. The organic layer was washed with brine and dried over MgSO_4 . Concentration gave a solid, which was purified by chromatography on silica gel eluting with mixtures of ethyl acetate and hexane.


- 5 LC retention time 5.28 minutes, $[\text{M}+\text{H}-\text{Boc}]^+$ 409.

Example 153



10

a)




- 15 The aldehyde (1 eq) from example 152 and 1-(3-Aminopropyl)-2-pipecoline (1.6 eq) were stirred at RT for 15min in 1,2-dichloroethane, then sodium triacetoxyborohydride (1.5) was added in one portion. After stirring for 16h at RT the mixture was concentrated *in vacuo* and the residue partitioned between dichloromethane and water. The organic layer was washed with brine and dried over MgSO_4 . Concentration gave a solid, which was purified
- 20 by chromatography on silica gel eluting with mixtures of dichloromethane and methanol. LC retention time 3.66 minutes, $[\text{M}+\text{H}]^+$ 649.

- b) The bis-Boc compound was treated with 1mL of 1/1 DCM/TFA and stirred at RT for 10 min. The volatiles were removed *in vacuo* to give the desired compound as the tris-trifluoroacetate.
- 25


LC retention time 2.92 minutes. $[\text{M}+\text{H}]^+$ 449.

Example 154



- 5 As example 153 using 1-(2-Aminoethyl)piperazine.
LC retention time 2.86 minutes, $[M+H]^+$ 422.


Example 155



10


- As example 153 using 1-(2-Aminoethyl)piperidine.
LC retention time 2.92 minutes, $[M+H]^+$ 421.

15 Example 156



- As example 153 using 4-(Aminomethyl)piperidine.
20 LC retention time 2.91 minutes, $[M+H]^+$ 407.


Example 157




- 5 To a mixture of the aldehyde (1 eq) of example 10, dichlorobis(triphenylphosphine)-palladium(II) (10 mol%) and 2N Na₂CO₃ was added 6-chloroindole-2-boronic acid (2 eq) in dry, degassed DME and the mixture heated at 80°C for 1h. After cooling to RT the solvent was removed *in vacuo* and the residue partitioned between ethyl acetate and water. The organic layer was washed with brine and dried over MgSO₄. Concentration gave a
- 10 solid, which was purified by chromatography on silica gel eluting with mixtures of ethyl acetate and hexane.
- LC retention time 5.16 minutes, [M+H-Boc]⁺ 429.

Example 158

15



a)



20

- The aldehyde (1 eq) from example 157 and 1-(3-Aminopropyl)-2-pipecoline (1.6 eq) were stirred at RT for 15min in 1,2-dichloroethane, then sodium triacetoxyborohydride (1.5) was added in one portion. After stirring for 16h at RT the mixture was concentrated *in vacuo*
- 25 and the residue partitioned between dichloromethane and water. The organic layer was


washed with brine and dried over MgSO_4 . Concentration gave a solid, which was purified by chromatography on silica gel eluting with mixtures of dichloromethane and methanol.

LC retention time 3.73 minutes, $[\text{M}+\text{H}]^+$ 669.

- 5 b) The bis-Boc compound was treated with 1mL of 1/1 DCM/TFA and stirred at RT for 10 min. The volatiles were removed *in vacuo* to give the desired compound as the tris-trifluoroacetate.

LC retention time 3.04 minutes. $[\text{M}+\text{H}]^+$ 469.


10 Example 159



As example 158 using 1-(2-Aminoethyl)piperazine.

- 15 LC retention time 2.97 minutes, $[\text{M}+\text{H}]^+$ 442.

Example 160




20

As example 158 using 1-(2-Aminoethyl)piperidine.

LC retention time 2.99 minutes. $[\text{M}+\text{H}]^+$ 441.

25


Example 161



5 As example 158 using 4-(Aminomethyl)piperidine.

LC retention time 3.00 minutes, $[M+H]^+$ 427.

Example 162



10


To a mixture of the aldehyde (1 eq) of example 10, dichlorobis(triphenylphosphine)-palladium(II) (10 mol%) and 2N Na_2CO_3 was added 5-fluoroindole-2-boronic acid (2 eq) in dry, degassed DME and the mixture heated at 80°C for 1h. After cooling to RT the

15 solvent was removed *in vacuo* and the residue partitioned between ethyl acetate and water. The organic layer was washed with brine and dried over MgSO_4 . Concentration gave a solid, which was purified by chromatography on silica gel eluting with mixtures of ethyl acetate and hexane.

LC retention time 5.31 minutes, $[M+H-\text{Boc}]^+$ 413.


20

Example 163



25


a)



- 5 The aldehyde (1 eq) from example 162 and N,N-Dimethylethylenediamine (1.6 eq) were stirred at RT for 15min in 1,2-dichloroethane, then sodium triacetoxyborohydride (1.5) was added in one portion. After stirring for 16h at RT the mixture was concentrated *in vacuo* and the residue partitioned between dichloromethane and water. The organic layer was washed with brine and dried over MgSO_4 . Concentration gave a solid, which was purified
10 by chromatography on silica gel eluting with mixtures of dichloromethane and methanol. LC retention time 3.59 minutes, $[\text{M}+\text{H}]^+$ 585.

- b) The bis-Boc compound was treated with 1mL of 1/1 DCM/TFA and stirred at RT for 10 min. The volatiles were removed *in vacuo* to give the desired compound as the tris-
15 trifluoroacetate. LC retention time 2.96 minutes, $[\text{M}+\text{H}]^+$ 385.

Example 164




20

As example 163 using 4-(Aminomethyl)piperidine.
LC retention time 2.89 minutes. $[\text{M}+\text{H}]^+$ 411.


25

Example 165



- 5 As example 163 using 1-(2-Aminoethyl)piperidine.
LC retention time 2.93 minutes, $[M+H]^+$ 425.

Example 166



10

- As example 163 using 1-(2-Aminoethyl)piperazine.
LC retention time 2.85 minutes, $[M+H]^+$ 426.

15 **Tat-Tar Binding Inhibition Assay***Principle of the assay*

- To measure the inhibition by the compound of RNA binding to ADP-1, the RNA is titrated
20 in the presence of a constant amount of fluorescent donor (fluorescein-ADP-1 peptide) and
compound as described in International Patent Application WO99/64625. The assay is
performed under competitive conditions, with a two fold excess of competitor RNA (a
fully base-paired TAR sequence) over fluorescein-ADP-1 peptide (the fluorescent donor).
The TAR RNA contains a 3' dabcyI group. The dabcyI group is a non-fluorescent acceptor
25 for energy transfer from fluorescein (the fluorescent donor). When ADP-1 and RNA bind,
the fluorescence signal from the fluorescein is quenched by the close proximity of the
DabcyI group. The presence of an inhibitory compound disrupts the ADP-1 RNA complex.

Complex disruption causes a decrease in energy transfer, which is observed as an increase in donor fluorescence intensity relative to a control (dabcyl) RNA-(fluorescein) ADP-1 binding reaction (in the absence of compound).

5 *Disruption of DABCYL-TAR-FAM-ADP-1 complex formation by compounds of the invention.*

Measurements were made in a 96-well plate reader (Wallac victor) with a fixed wavelength of 490nm and emission at 535 nm. I_0 was determined by an initial measurement of a 95 μ L solution of 10nM Fluorescein-ADP-1 in the presence of 50mM Tris.HCl pH7.5, 80mM KCl, 1% DMSO, 0.01% Triton X-100, 5 μ g/mL BSA, 20nM competitor RNA in the presence 1 μ M compound. I was then measured following the addition of 5 μ L of a 20 X DABCYL-TAR RNA stock solution.

Example No.	Ki(μ M)	Example No.	Ki(μ M)	Example No.	Ki(μ M)
2	<10	40	<10	87	<1
3	<10	42	<1	88	<10
4	<1	43	<1	89	<10
5	<10	44	<1	90	<50
6	<1	45	<50	91	<1
7	<10	46	<1	92	<10
8	<1	48	<10	93	<1
9	<1	49	<10	94	<1
11	<10	50	<10	95	<10
12	<10	51	<1	96	<1
13	<1	52	<50	98	<1
14	<1	54	<1	99	<10
15	<1	55	<50	100	<10
16	<10	56	<1	107	<1
17	<10	58	<10	108	<1
18	<1	59	<50	109	<10
19	<1	64	<1	110	<1

21	<1	65	<10	111	<1
22	<1	66	<1	112	<1
23	<1	68	<1	113	<1
24	<1	69	<10	116	<10
26	<1	70	<1	117	<10
27	<1	71	<10	118	<10
28	<50	74	<50	119	<50
29	<10	75	<50	120	<10
30	<10	77	<50	121	<10
31	<10	78	<50	126	<10
32	<10	79	<50	127	<10
33	<1	80	<10	129	<10
34	<10	81	<50	136	<10
35	<10	82	<100	137	<10
36	<1	83	<50	139	<1
37	<1	85	<1	140	<1
38	<10	72	<100		
39	<10	86	<10		

In vitro Translation assay

Compounds of the present invention showed inhibitory activities in *in vitro* translation assays utilizing *E. coli* extracts. The plasmid pBestLuc, which contained the gene for firefly luciferase downstream of an *E. coli* promoter and a ribosome binding site was used as a template. The activity of the firefly luciferase enzyme resulted in a strong luminescent signal. The luminescence generated was a direct measurement of protein expression and of translation efficiency.

10

Translation reactions in the presence of compound were started by mixing a translation premix that contained Mg^{2+} , plasmid template, amino acids, nucleotidetriphosphates, phosphocreatine, creatine phosphokinase and folinic acid with the S30 extract that contains RNA polymerase, ribosomes and translation factors (prepared from *E. coli* MRE600 cells) followed by incubation at 37°C. The activity of the translated luciferase protein was

15

measured by adding an aliquot of the translation reaction to the non-luminescent substrate luciferin and the luminescence measured. The luminescence was quantified in a luminescence plate reader (Wallac Victor). Compounds were assayed 3-5 times over a range of concentrations and an IC_{50} calculated.

5

Example No.	8	13	14	15	51	85	98	110	112	113
IC_{50} (μM)	26.7	16.7	17.8	18.1	16.8	27.2	7.1	18.2	5.6	8.5

In vivo antibacterial assay

The *in vivo* therapeutic efficacy of the compounds of the invention is measured by intramuscular injection to mice experimentally infected with a pathogenic gram positive or gram negative bacterium (e.g. methicillin-resistant *Staphylococcus aureus* (MRSA), *Clostridium difficile*, *Klebsiella pneumoniae*, *Escherichia coli*, *Haemophilus influenzae*, etc.). As an example, MRSA strain A27223 can be used.

MRSA strain A27223 is prepared for experimental infection by growth on two large Brain Heart Infusion Agar plates. On each plate, 0.5 ml of frozen stock culture is plated out. Plates are then incubated for 18 hours at 30°C. The next day each plate is washed with 20 ml of Brain Heart Infusion Broth and then pooled together. A microscopic direct count of microorganisms is done using a 1:1000 dilution of plate wash. After a direct count is obtained, the number of organisms per milliliter is calculated. The count is adjusted to the desired amount of inoculum by diluting in 4% hog mucin. The desired challenge (amount of organisms given to mice) is 2.4×10^8 cfu/0.5 ml/mouse for MRSA strain A27223. The mice are infected intraperitoneally with 0.5 ml of challenge. Ten non-treated infected mice are used as controls. Mice used are adult male ICR mice. The average weight of the animals should range from 20 to about 26 grams.

Compounds are generally tested at 4 dose levels (e.g. 25, 6.25, 1.56 and 0.39 mg/kg) and prepared in 5% cremophor, unless otherwise specified. When MRSA A27223 is the challenging microorganism, vancomycin is used as the control compound, and is dosed at

6.25, 1.56, 0.39 and 0.098 mg/kg. It is prepared in 0.1M phosphate buffer. There are generally five infected mice per dose level, and they are treated with 0.2 ml of the test compound, preferably by intramuscular injection. Treatment begins 15 minutes and 2 hours post-infection.

5

A PD₅₀ (protective dose-50, the dose of drug given which protects 50% mice from mortality) runs for 5 days. During this time, mortality of mice is checked every day and deaths are recorded. The cumulative mortality at each dose level is used to calculate a PD₅₀ value for each compound. Surviving mice are sacrificed at the end of day 5 by CO₂ inhalation. The actual calculation of PD₅₀ is performed with a computer program using the Spearman-Kärber procedure.

10

A compound according to the invention is effective for the treatment of bacterial infection if it has a PD₅₀ of about 100 mg/kg or less.

15

In vivo antiviral assay

The *in vivo* therapeutic efficacy of the compounds of the invention is measured by conventional *in vivo* antiviral assays including, but not limited to, that described in Letvin,

20


N.L., Daniel, M.D., Sehgal, P.K., Desrosiers, R.C., Hunt, R.D., Waldron, L.M., MacKey, J.J., Schmidt, D.K., Chalifoux, L.V. and King, N.W. Introduction of AIDS-like disease in macaque monkeys with T-cell tropic retrovirus STLV-III, *Science*, 1985, 230, 71-73, which is incorporated herein by reference.

25

Variations, modifications, and other implementations of what is described herein will occur to those of ordinary skill in the art without departing from the spirit and scope of the invention as claimed. Accordingly, the invention is to be defined not by the preceding illustrative description but instead by the spirit and scope of the following claims.

CLAIMS:

1. A compound of the formula



- 5 wherein

Ar is an aryl group,

X¹ is selected from O, S, SO, SO₂, and NR,

X² is selected from O, S, SO, SO₂, NR and CR₂,

X³ is CR₂,

- 10 Y¹ and Y² are independently selected from C₁₋₁₂ alkylene, C₄₋₁₂ arylene, C₄₋₁₆ aralkylene, CO(C₁₋₁₂ alkylene), CO(C₄₋₁₂ arylene) and CO(C₄₋₁₆ aralkylene) groups,

A and B are independently selected from groups comprising a group selected from:

- 15 amine (-NR₂), amide (-CONR₂), amidine (-C(=NR)NR₂), thioamide (-CSNR₂), oxime (=NOR), hydroxylamine (-NHOR), hydroxamic acid (-CONROR), hydrazine (-NRNR₂), hydrazone (=NNR₂), sulphonamide (-SO₂NR₂), sulphinamide (-SONR₂), sulfoximine (-SO(=NR)-), urea (-NRCONR₂), guanidine (-NRC(=NR)NR₂), and aromatic and non-aromatic nitrogen heterocyclic groups,

each R is independently selected from H, C₁₋₁₂ alkyl and C₃₋₁₂ aryl, or any two R groups

- 20 may together comprise a C₁₋₆ alkylene chain,
or a pharmaceutically acceptable derivative thereof.

2. A compound according to claim 1 wherein X¹ is O.


- 25 3. A compound according to claim 1 or 2 wherein X² is NR.

4. A compound according to claim 3 wherein X² is NH.

5. A compound according to any preceding claim wherein X³ is CH₂.

6. A compound according to any preceding claim wherein Ar is a monocyclic or a fused bicyclic aromatic or heteroaromatic group.

7. A compound according to claim 6 wherein Ar is a fused bicyclic heteroaromatic group
5 of the formula



wherein X^4 is NH, S or O.

8. A compound according to claim 7 wherein X^4 is NH.

10

9. A compound according to any preceding claim wherein Y^1 comprises a C_{1-5} alkylene group.

15

10. A compound according to any preceding claim wherein Y^2 comprises a C_{1-5} alkylene group.

20

11. A compound according to any preceding claim wherein A is a group comprising a group selected from amine, amidine, guanidine, and aromatic and non-aromatic nitrogen heterocyclic groups.

12. A compound according to any preceding claim wherein B is a group comprising a group selected from amine, amidine, guanidine, and aromatic and non-aromatic nitrogen heterocyclic groups.

25 13. A compound according to any one of claims 1 to 12 for use in therapy.

14. Use of a compound according to any one of claims 1 to 12 in the manufacture of a medicament for use in the treatment of viral infection or bacterial infection.

15. A method of treating viral infection or bacterial infection comprising administering to a patient in need of such treatment an effective dose of a compound according to any one of claims 1 to 12.

5 16. A pharmaceutical composition comprising a compound according to any one of claims 1 to 12 in combination with a pharmaceutically acceptable excipient.

17. Use of a compound according to any one of claims 1 to 12 to inhibit the binding of Tat to Tar or to inhibit bacterial protein translation.

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/GB 01/00362

A. CLASSIFICATION OF SUBJECT MATTER					
IPC 7	C07D211/26	C07D243/08	C07D333/16	C07D333/56	C07D307/80
	C07D307/85	A61K31/155	A61K31/381	A61K31/343	A61K31/404
	A61K31/4164	A61K31/445	A61K31/4025	A61K31/506	A61K31/501
According to International Patent Classification (IPC) or to both national classification and IPC					
B. FIELDS SEARCHED					
Minimum documentation searched: (classification system followed by classification symbols)					
IPC 7	C07D	C07C	A61K	A61P	
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched					
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)					
CHEM ABS Data, EPO-Internal, PAJ, WPI Data					
C. DOCUMENTS CONSIDERED TO BE RELEVANT					
Category *	Citation of document, with indication, where appropriate, of the relevant passages				Relevant to claim No.
A	EP 0 580 402 A (SQUIBB & SONS INC) 26 January 1994 (1994-01-26) abstract				1, 13-17
A	--- DATABASE CHEMABS 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; CHANG, XINGRUO ET AL: "Studies on the chemical constituents of Glycyrrhiza. I. Isolation and structural elucidation of the antibacterial constituent, licobenzofuran" retrieved from STN Database accession no. 97:20701 XP002147477 abstract; figures & ZHONGCAOYAO (1981), 12(12), 530, --- -/--				1, 13-15
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.					
* Special categories of cited documents: *A* document defining the general state of the art which is not considered to be of particular relevance *E* earlier document but published on or after the international filing date *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the international filing date but later than the priority date claimed *I* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. *Z* document member of the same patent family					
Date of the actual completion of the international search			Date of mailing of the international search report		
26 April 2001			11/05/2001		
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040. Tx. 31 651 epo nl. Fax: (+31-70) 340-3016			Authorized officer Paisdor, B		

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB 01/00362

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/5513 C07D209/12 C07D209/08 C07D213/30 C07D311/22
C07D215/14 C07D239/52 C07D209/42 C07D209/86 C07D277/24
C07C279/12 C07D295/20 C07D409/12 C07D401/12 C07D403/12

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	BARRON, D. I. ET AL: "Synthesis and antiinflammatory activity of 4-(p-biphenyl)-3- hydroxybutyric acid and related compounds" J. MED. CHEM. (1968), 11(6), 1139-44, XP002147476 page 1140; examples 23-39; table I ---	1,13
A	GB 1 171 251 A (BRITISH DRUG HOUSES LTD) 19 November 1969 (1969-11-19) page 3 -page 14; examples page 1, line 7 - line 9; claim 1 --- -/-	1,13



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

26 April 2001

Date of mailing of the international search report

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.
Fax: (+31-70) 340-3016

Authorized officer

Paisdor, 8

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 01/00362

A. CLASSIFICATION OF SUBJECT MATTER
 IPC 7 A61P31/12 A61P31/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	PATENT ABSTRACTS OF JAPAN vol. 007, no. 034 (C-150), 10 February 1983 (1983-02-10) -& JP 57 188579 A (TAKEDA YAKUHIIN KOGYO KK;OTHERS: 01), 19 November 1982 (1982-11-19) abstract page 622 -page 623; tables --- -/--	1,13

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

26 April 2001

Date of mailing of the international search report

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Paisdor, B

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 01/00362

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	KUMAR S ET AL: "P-(CYCLOPENTENYL)PHENOL DERIVATIVES AS POTENTIAL BIODYNAMIC AGENTS" INDIAN JOURNAL OF CHEMISTRY, SECTION B: ORGANIC, INCL. MEDICINAL, IN, PUBLICATIONS & INFORMATIONS DIRECTORATE, NEW DELHI, vol. 25B, 1968, pages 106-110, XP000909818 ISSN: 0019-5103 page 106, column 2, paragraph 3 page 107; examples 4-6	1,13
A	WO 96 11902 A (ZENECA LTD ;BREAU LT GLORIA ANN (GB); OLDFIELD JOHN (GB); TUCKER HO) 25 April 1996 (1996-04-25) claims 1,13	1,13
A	GB 2 086 386 A (MAY & BAKER LTD) 12 May 1982 (1982-05-12) claims	1,13
A	WO 97 48674 A (ORTHO PHARMA CORP) 24 December 1997 (1997-12-24) abstract; claims	1,13-17
A	EP 0 574 808 A (THOMAE GMBH DR K) 22 December 1993 (1993-12-22) abstract; claims	1,13-17
A	EP 0 496 378 A (THOMAE GMBH DR K) 29 July 1992 (1992-07-29) abstract; claims	1,13-17
A	EP 0 144 892 A (HOECHST AG) 19 June 1985 (1985-06-19) claims	1,13-17
A	US 4 129 572 A (PIGEROL CHARLES ET AL) 12 December 1978 (1978-12-12) abstract; claim 1 column 5 -column 8; examples	1

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 01/00362

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0580402 A	26-01-1994	US 5559256 A	24-09-1996
		AU 677194 B	17-04-1997
		AU 4165993 A	27-01-1994
		CA 2100894 A	21-01-1994
		CN 1085546 A	20-04-1994
		FI 933277 A	21-01-1994
		HU 67090 A	30-01-1995
		HU 9500679 A	28-11-1995
		JP 6206857 A	26-07-1994
		NO 932620 A	21-01-1994
		US 5760036 A	02-06-1998
		US 5776933 A	07-07-1998
		BG 97961 A	31-03-1995
		CZ 9301437 A	15-02-1995
		MX 9304351 A	28-02-1994
		PL 299738 A	05-04-1994
		SK 76093 A	05-10-1994
		ZA 9305243 A	17-02-1994
GB 1171251 A	19-11-1969	AT 281038 B	11-05-1970
		CH 496005 A	15-09-1970
		DE 1770805 A	13-01-1972
		DK 120998 B	16-08-1971
		ES 355739 A	16-12-1969
		FR 1583999 A	12-12-1969
		IE 32152 B	02-05-1973
		IL 30313 A	30-08-1972
		NL 6809422 A	07-01-1969
		SE 352891 B	15-01-1973
		US 3732229 A	08-05-1973
JP 57188579 A	19-11-1982	JP 1014915 B	14-03-1989
		JP 1531952 C	24-11-1989
		DK 215082 A	16-11-1982
		ES 512185 D	01-02-1983
		ES 8303374 A	01-05-1983
WO 9611902 A	25-04-1996	AU 3616295 A	06-05-1996
		DE 69514087 D	27-01-2000
		DE 69514087 T	27-04-2000
		EP 0733033 A	25-09-1996
		JP 9511529 T	18-11-1997
		US 5811459 A	22-09-1998
		ZA 9508622 A	12-04-1996
GB 2086386 A	12-05-1982	AU 547334 B	17-10-1985
		AU 7674581 A	06-05-1982
		BE 890874 A	26-04-1982
		CH 650776 A	15-08-1985
		DE 3142559 A	02-09-1982
		DK 471181 A, B,	28-04-1982
		EG 15900 A	30-10-1987
		FR 2492821 A	30-04-1982
		GR 75366 A	13-07-1984
		IE 51670 B	04-02-1987
		IL 64123 A	28-02-1986
		IT 1139562 B	24-09-1986
		JP 57099573 A	21-06-1982

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 01/00362

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB 2086386 A		KE 3690 A	20-02-1987
		LU 83713 A	07-05-1982
		MY 55685 A	31-12-1985
		NL 8104766 A	17-05-1982
		OA 7362 A	30-06-1984
		PH 17779 A	11-12-1984
		US 4374136 A	15-02-1983
		ZA 8107298 A	29-09-1982
		ZW 25681 A	13-01-1982
WO 9748674 A	24-12-1997	US 5773469 A	30-06-1998
		AU 3480497 A	07-01-1998
		ZA 9705331 A	17-12-1998
EP 0574808 A	22-12-1993	DE 4219158 A	16-12-1993
		AU 4120193 A	23-12-1993
		CA 2098158 A	12-12-1993
		CN 1080917 A	19-01-1994
		FI 932649 A	12-12-1993
		JP 6073038 A	15-03-1994
		MX 9303466 A	31-01-1994
		NO 932120 A	13-12-1993
		PL 299260 A	07-03-1994
		ZA 9304090 A	11-12-1994
EP 0496378 A	29-07-1992	DE 4102024 A	30-07-1992
		AT 128120 T	15-10-1995
		AU 648379 B	21-04-1994
		AU 1040392 A	30-07-1992
		CA 2059857 A	25-07-1992
		DE 59203704 D	26-10-1995
		DK 496378 T	05-02-1996
		ES 2079694 T	16-01-1996
		GR 3017802 T	31-01-1996
		HU 60462 A	28-09-1992
		IE 920199 A	29-07-1992
		JP 4334351 A	20-11-1992
		MX 9200269 A	01-07-1992
		NO 177852 B	28-08-1995
		NZ 241355 A	26-07-1994
		US 5597825 A	28-01-1997
		US 5736559 A	07-04-1998
		US 5922763 A	13-07-1999
		ZA 9200464 A	23-07-1993
EP 0144892 A	19-06-1985	DE 3343815 A	13-06-1985
		AT 38380 T	15-11-1988
		DE 3474957 D	08-12-1988
		DK 572084 A	04-06-1985
		ES 538124 D	16-03-1986
		ES 8605499 A	01-09-1986
		GR 81123 A	26-03-1985
		JP 60136556 A	20-07-1985
		PT 79602 A, B	01-01-1985
US 4129572 A	12-12-1978	FR 2358395 A	10-02-1978
		BE 856620 A	09-01-1978
		DE 2731456 A	19-01-1978

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 01/00362

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4129572 A		GB 1541957 A	14-03-1979
		JP 53015368 A	13-02-1978
		NL 7707793 A	17-01-1978
		SE 422459 B	08-03-1982
		SE 7708186 A	16-01-1978
<hr/>			